

Prevalence and Associated Factors of Tardive Dyskinesia Among Psychiatric Patients on First-Generation Antipsychotics at Jimma University Specialized Hospital, Psychiatric Clinic, Ethiopia: Institution Based on A Cross-Sectional Study

Tilahun Abdeta^{1*}, Daniel Tolessa², Wondale Tsega³

¹Department of Psychiatry, School of Nursing and Midwifery, College of health and medical sciences, Haramaya University, Dire Dawa, Ethiopia

²Department of Psychiatry, Adama Hospital medical College, Adama, Ethiopia

³Department of medicine, Adama Hospital medical college, Adama, Ethiopia

***Corresponding Author:** Dr. Tilahun Abdeta, Department of Psychiatry, School of Nursing and Midwifery, College of health and medical sciences, Haramaya University, Dire Dawa, Ethiopia, E-mail: tilahunabdeta@gmail.com

Received: 18 July 2019; **Accepted:** 09 August 2019; **Published:** 12 August 2019

Abstract

Background: Tardive Dyskinesia is described as a movement disorder that can be caused by the prolonged use of antipsychotics. It is a permanent condition that affects quality of life. However, studies on TD among psychiatric patients in Ethiopia are limited. The objective of this study was to determine the prevalence and associated factors of Tardive dyskinesia among psychiatric patients on first-generation antipsychotics at Jimma University specialized hospital (JUSH) psychiatric clinic, Jimma, South west Ethiopia.

Method: Hospital based cross-sectional study was conducted using Abnormal Involuntary Movement Scale (AIMS) and Schooler and Kane research criteria to identify cases of first-generation antipsychotic-induced TD among 151 psychiatric outpatients on first-generation antipsychotics at Jimma University Specialized Hospital psychiatric clinic. Simple random sampling method was employed to select subjects from a total of 912 patients on follow-up and logistic regression analysis was employed to identify associated factors.

Result: In our study the prevalence of first-generation antipsychotic induced TD was 14.6% (95%CI: 10.76%, 18.4%). In the final model multivariate logistic regression analysis revealed that age (AOR 4.5, 95%CI: 9.7, 20.4), current smoking cigarettes (AOR 1.4, 95%CI: 2.6, 7.8) and chlorpromazine equivalent dose (AOR 6.5, 95%CI: 2.6, 26.8) had statistically significant association with first-generation antipsychotic induced TD.

Conclusion: Considerable number of psychiatric patients were suffered from first-generation antipsychotic induced TD. Age and current smoking cigarettes and chlorpromazine equivalent dose had significant association with first-generation antipsychotic induced TD. Therefore, stakeholders should consider and appropriately address these statistically identified associated factors in order to reduce antipsychotic induced TD.

Keywords: Prevalence; Associated factors; TD- Tardive Dyskinesia; First-generation antipsychotics; JUSH- Jimma University Specialized Hospital

Abbreviations: AIMS- Abnormal Involuntary Movement Scale; AOR- Adjusted Odds Ratio; COR- Crude Odds Ratio; JUSH- Jimma University Specialized Hospital; OPD- Outpatient Department; TD- Tardive Dyskinesia

1. Background

First-generation antipsychotics-induced movement disorders constitute a worldwide problem in the treatment of schizophrenia and related psychotic disorders because of the limited affordability of relatively safe second-generation antipsychotics [1]. Tardive dyskinesia (TD) is one of a group of side effects called antipsychotics induced movement disorders. TD is perhaps the most severe of these side effects and it is one of the most disturbing potential side effects [2]. Tardive Dyskinesia is a movement disorder that occurs over months, years and even decades after started antipsychotic medications [2]. The annualized TD incidence across first-generation antipsychotics arms was 6.5% (95% CI: 5.3-7.8%) vs. 2.6% (95% CI: 2.0-3.1%) across second-generation antipsychotics arms. TD risk and annualized rates were lower with second generation antipsychotics compared to first-generation antipsychotics (RR=0.47, 95% CI: 0.39-0.57, $p < 0.0001$, $k=28$; RaR=0.35, 95% CI: 0.28-0.45, $p < 0.0001$, number-needed-to-treat, NNT=20) [3-5]. First-generation antipsychotic drugs are efficacious in the treatment of acute psychosis, chronic psychotic conditions and other psychiatric disorders [6]. Drugs in this class have comparable efficacy in the treatment of psychosis and include high potent drugs (haloperidol, thiothixene, fluphenazine, Trifluoperazine, perphenazine, loxapine, droperidol & pimozide) and low potent drugs (chlorpromazine, thioridazine & mesoridazine) [6]. Tardive Dyskinesia is characterized by a condition of potentially irreversible abnormal involuntary movements of different muscles within the body and can occur in the tongue, lips or jaw (e.g., facial grimacing), or consist of purposeless movements of arms, legs, fingers and toes. In some severe cases, TD can include swaying movements of the trunk or hips or affect the muscles associated with breathing and can be quite embarrassing and depending on its severity can be disabling as well. It results in behavioral disturbances, non-adherence, and exacerbation of psychosis [2].

Literatures indicated various prevalence of TD. A meta-analysis study revealed that 6.5 % of patients who were treated with first-generation and predominantly high-potency antipsychotic agents, such as haloperidol or fluphenazine met diagnostic criteria for TD [3]. The cumulative incidence of TD after antipsychotic drug exposure of 1, 5, and 10 years was 5, 27 and 43 percent, respectively [7]. Studies from non-western countries, including Africa are sparse. The study in African Caribbean patients on long-term antipsychotic treatment, showed that in patients with a mean antipsychotic use of approximately 18 years, the yearly incidence rate of TD was 10.2% (95% CI = 7.7 to 13.5) [8]. Several factors in relation with drug treatment have been identified as contributing to the risk of developing TD includes: age, psychiatric diagnosis, gender, organic brain damage, negative symptoms in patients

with schizophrenia, antipsychotic dose, duration of treatment, the number of drug free intervals and the history of acute extra-pyramidal side effects [1, 9].

In spite mental health problem was included in national health policy of Ethiopia, interventions for them are limited. One of the main reasons behind is lack of data on the extent of the problem. Search for literatures revealed, there were very limited published data that investigated the prevalence and associated factors of first-generation antipsychotics induced TD in Ethiopia. So, our study result will add valuable points to or strengthens the knowledge about the prevalence and associated factors of tardive dyskinesia in second world countries including Ethiopia. Therefore, the objective of this study was to determine the prevalence and associated factors of Tardive dyskinesia among psychiatric patients on first-generation antipsychotics at Jimma University specialized hospital (JUSH) psychiatric clinic, Jimma, South west Ethiopia.

2. Materials and Methods

2.1 Study area and period

The study was conducted from Jun 21 to July 5, 2016 at Jimma University Specialized Hospital (JUSH), Psychiatric clinic to assess the prevalence of TD and associated factors. Jimma University Specialized Hospital is one of the Hospitals in Ethiopia which is found in Jimma city, located 345 km southwest of Addis Ababa. It is a referral center for the southwestern part of Ethiopia, and currently, the only teaching and referral hospital in the southwestern part of the country. Jimma University Specialized Hospital Psychiatric clinic gives both inpatient and outpatient service for psychiatric patients mostly coming from Jimma area as well as patients referred from other health institutions in the southwestern region of the country. Even if majority of the patients come from Jimma area, the hospital give service for patients come from up to 300 km radius from Jimma town in the southwest part of Ethiopia. Generally, it gives services for more than 30 million populations, which served a huge bulk of the community next to Paulos Hospital in Addis Ababa Ethiopia concerning Psychiatry health facility.

2.2 Study design

Hospital based Cross sectional study design was employed.

2.3 Source population

All psychiatric patients on first-generation antipsychotics at Jimma University Specialized Hospital, Psychiatric clinic.

2.4 Study population

All Psychiatric patients on first-generation antipsychotics who were attending follow up at Jimma University Specialized Hospital, Psychiatric clinic outpatient department (OPD) during the study period and fulfill the inclusion criteria.

2.4.1 Inclusion criteria: All psychiatric patients who were on first-generation anti-psychotic drugs treatment at least for 6 months.

2.4.2 Exclusion criteria: Psychiatric patients whose abnormal involuntary movements were induced by other reasons like infections and seizure were excluded;

2.5 Sample size determination and Sampling technique

Using single population proportion formula and taking the prevalence of antipsychotic induced movement disorders (Tardive Dyskinesia) from the study conducted at Amanuel mental specialized hospital, Addis Ababa, Ethiopia [1] which was 11%, with 95% confidence level and marginal error of 5% the final sample size was 151 individuals. Simple random sampling technique was used to select the study participants. From a total of 912 psychiatric patients who were taking first-generation antipsychotics and on follow-up at Jimma University specialized hospital, psychiatric clinic, we collected their card number from card room and we selected the total study participants (151 individuals) using simple random sampling technique by lottery method.

2.6 Study variables

2.6.1 Dependent variable: The presence of Tardive Dyskinesia (yes/no).

2.6.2 Independent Variables: The independent variables included in this study were: Age, Sex, Religion, Ethnicity, Occupational status, Educational status, Current Alcohol use, Current Khat use, Current smoking Cigarette, Current Caffeine use, Diagnosis, Duration of psychiatric disorder, Drug used and Duration of treatment.

2.7 Data collection method and tools

Data were collected by four (4) experienced psychiatric nurses using interviewer administered structured questionnaire and research criteria suggested by Schooler and Kane. The questionnaire used was Abnormal Involuntary Movement Scale (AIMS) which is a rating scale designed in the 1970s to measure involuntary movements known as tardive dyskinesia (TD) [10]. The AIMS test has a total of twelve items rating involuntary movements of various areas of the patient's body. These items are rated on a five-point scale of severity from 0–4. The scale is rated from 0 (none), 1 (minimal), 2 (mild), 3 (moderate), 4 (severe) and the entire test performed within ten (10) minutes for each patient. For the diagnosis of Tardive dyskinesia (TD), research criteria suggested by Schooler and Kane [11] were used and they are as follows: 1. At least 3 months of total exposure to ant-dopaminergic medicinal products, 2. Exclusion of other reasons for involuntary movements, and 3. A minimum global rating of “mild”(i.e. 2 or more on AIMS) was used.

Before data collection time the data collectors were trained by principal investigators on the objectives of the study and how to identify Tardive Dyskinesia in accordance with Abnormal Involuntary Movement Scale (AIMS) for Tardive dyskinesia.

2.8 Data analysis and processing

The Data were analyzed using SPSS software program version 20. In order to determine association between TD and independent variables, we employed Logistic regression analysis. Characteristics of study population were described by descriptive statistics. Odds ratio with 95% confidence interval was calculated to assess the association and statistical significance. Those variables which were found to be significant in the bivariate analysis with a p-

value <0.05 were retained for further multivariate logistic regression analysis to control for potential confounders and to predict independent factors associated with TD.

2.9 Data quality assurance

To maintain the quality of the data, data collectors were trained on how to approach respondents, how to collect and fill all the necessary data and on different variables which are used in the study. Meanwhile, the data collectors were supervised by the principal investigator and One week before the actual survey pretest was carried out among patients with similar socio-demographic characteristic and using similar rating scale which was, Abnormal Involuntary Movement Scale (AIMS) for Tardive dyskinesia. The pretest was conducted at Jimma University Specialized Hospital, psychiatric clinic on 5% of the total sample size and necessary modification was done before being applied on the study subject. Also, the result of pre-test was not included in the final result of the study. Concerning inter rater reliability of data collectors, only 4 experienced psychiatric nurses collected the data under close supervision of the principal investigator.

2.10 Operational definitions

2.10.1 Tardive dyskinesia (TD): In this study TD is defined as patients who scored a minimum global rating of “mild”(i.e. 2 or more on AIMS) [11].

2.10.2 Current alcohol, Khat, cigarettes and caffeine use: Defined as the proportion of individuals who used these substances at least once during the last 1 month preceding the study [12].

2.11 Ethical clearance

Ethical clearance was obtained from the Research Ethical Review Board of Jimma University, institute of health, and the study was performed according to the declaration of Helsinki. The respondents were informed about their right to refuse or agree to participate in the study and written informed consent was obtained from every participant before the start of the data collection.

3. Results

3.1 Socio-demographic characteristics of respondents

All 151 participants involved in this study agreed providing response rate of 100%, 70.2% were males and 49% were found in the age range of less than 30 years. The majority of participants (76.8%) were Oromo by their ethnicity, 67.5% were Muslims by their religion, 42.4% were jobless and the 42.4% of respondents' educational status were 1-6 grades (Table 1).

Socio-demographic characteristics		Count (N)	Percentage (%)
Educational status	Can't read and write	20	13.2%
	1-6 grades	64	42.4%
	7-12 grades	43	28.5%
	>12 grades	24	15.9%

Ethnicity	Oromo	116	76.8%
	Amhara	17	11.3%
	Kaffa	10	6.6%
	Others	8	5.3%
Religion	Orthodox	33	21.9%
	Muslim	102	67.5%
	Protestant	16	10.6%
Occupation	Has job	87	57.6%

Table 1: Socio-demographic characteristics of psychiatric patients on first-generation antipsychotics at JUSH, psychiatric clinic, Jimma, Ethiopia, 2016.

Majority of the total participants were found within the age category of <30 years (Figure 1) and 79.5% had the diagnosis of schizophrenia and 47% had gone 5-9 years with their psychiatric illness. The 38.4% of patients were using chlorpromazine and 49% of participants took their treatment drugs for more than 5 years and the majority (74.8%) of the patients reported as they were using caffeine (Table 2).

Clinical characteristics		Count (N)	Percentage (%)
Diagnosis of patient	Schizophrenia	120	79.5%
	Others*	31	20.5%
Duration of psychiatric disorder	<5 year	61	40.4%
	5-9 year	71	47.0%
	>= 10 year	19	12.6%
Types of Drugs used	Haloperidol	48	31.8%
	Chlorpromazine	58	38.4%
	Thioridazine and Trifluoperazine	45	29.8%
Duration of treatment	3 month - < 6 months	6	4.0%
	6 month – 5 year	71	47.0%
	>= 5 year	74	49.0%
Chlorpromazine equivalent dose (mg/day)	50-<100	13	8.6
	100-<400	112	74.2
	>=400	26	17.2
Current Alcohol use	Yes	6	4.0%
Current Caffeine use	Yes	113	74.8%
Current Smoking cigarette	Yes	12	7.9%
Current Khat use	Yes	70	46.4%

Others* (Bipolar disorder and Major Depressive Disorder with psychotic features)

Table 2: Clinical and current substance use characteristics of psychiatric patients on first-generation anti-psychotics at JUSH, psychiatric clinic, Jimma, Ethiopia, 2016.




Figure 1: Age and sex of psychiatric patients who were on first-generation antipsychotics at JUSH psychiatric clinic, 2016.

3.2 Prevalence of first-generation antipsychotics induced Tardive Dyskinesia (TD)

In the current study the prevalence of first-generation antipsychotic induced Tardive dyskinesia among psychiatric patients on first-generation antipsychotics at Jimma University specialized hospital, psychiatric clinic, Ethiopia was 14.6% (95%CI: 10.76%, 18.4%).

3.3 Factors associated with first-generation antipsychotic induced Tardive Dyskinesia

In order to identify the presence of association between first-generation antipsychotic induced Tardive dyskinesia and independent variables we performed bivariate and multivariate logistic regression analyses. In bivariate logistic regression analysis patients' age, religion, chlorpromazine equivalent dose, current alcohol use, and current smoking cigarettes were found having significant association with first-generation antipsychotic induced tardive dyskinesia and they were candidates for final model of multivariate logistic regression analysis. The final model of multivariate logistic regression analysis showed that age, chlorpromazine equivalent and current smoking cigarettes had statistically significant association with first-generation antipsychotics induced Tardive dyskinesia. The final model of multivariate logistic regression analysis revealed that patients who had smoked cigarettes at least once during the last one month were 1.4 times more likely to develop TD as compared to those who were not smoking cigarettes during the last one month (AOR 1.4, 95%CI: 2.6, 7.8) and patients who were in the age category ≥ 45 year were more than four times more likely to develop TD as compared to those who were found in the age category of <30 year (AOR 4.5, 95%CI: 9.7, 20.4). Patients who have been on chlorpromazine equivalent dose range ≥ 400 mg/day were 6.5 times more likely to develop TD compared to patients who have been on chlorpromazine equivalent dose range 50-<100 mg/day (AOR 6.5, 95%CI: 2.6, 26.8) (Table 3).

Variables	Antipsychotic induced TD		p-value	COR (95%CI)	p-value	AOR (95%CI)
	Yes	No				
Age						
<30	7	68	Reference	Reference	Reference	Reference
30-44	11	48	0.15	2.4(0.7-7.8)	0.2	2.6(0.6-11.3)
≥45	6	11	0.01*	5.3(1.5-18.7)	0.04*	4.5(9.7-20.4)
Sex						
Male	15	91	Reference	Reference	Reference	Reference
Female	9	36	0.37	0.66(0.26-1.6)		
Educational status						
Can't read and write	4	16	0.27	0.36(0.1-2.2)		
1-6 grades	8	56	0.58	0.64(0.12-3.2)		
7-12 grades	10	33	0.14	0.3(0.1-1.5)		
>12 grades	2	22	Reference	Reference	Reference	Reference
Occupation						
Has job	12	75	Reference	Reference	Reference	Reference
No job	12	52	0.41	0.69(0.29-1.7)		
Ethnicity						
Oromo	15	101	Reference	Reference	Reference	Reference
Amhara	4	13	0.25	0.48(0.14-1.7)		
Kaffa	3	7	0.15	0.35(0.1-1.5)		
Others	2	6	0.35	0.45(0.1-2.4)		
Religion						
Orthodox	10	23	0.02*	3.3(1.3-8.5)	0.06	3.6(0.95-14.0)
Muslim	12	90	Reference	Reference	Reference	Reference
Protestant	2	14	0.19	3.0(0.6-15.9)	0.13	5.1(0.62-42.3)
Diagnosis of patient						
Schizophrenia	17	103	Reference	Reference	Reference	Reference
Others	7	24	0.26	0.57(0.21-1.5)		
Duration of psychiatric disorder						
<5 year	9	52	Reference	Reference	Reference	Reference
5-9 year	11	60	0.91	0.94(0.36-2.5)		
≥ 10 year	4	15	0.52	0.65(0.17-2.4)		
Drug used						
Haloperidol	10	38	0.51	0.7(0.24-2.0)		
Chlorpromazine	7	51	0.61	1.3(0.43-4.15)		
Thioridazine and	7	38	Reference	Reference	Reference	Reference

Trifluoperazine						
Duration of treatment						
3months -< 6 months	1	5	Reference	Reference	Reference	Reference
6 month -5year	9	62	0.78	1.4(0.1-13.12)		
>=5 year	14	60	0.89	0.8(0.09-7.9)		
Chlorpromazine equivalent dose (mg/day)						
50-<100	2	11	Reference	Reference	Reference	Reference
100-<400	8	104	0.03*	6.4(11.8-34.8)	0.06	4.3(0.92-16.9)
>=400	14	12	0.001*	10.2(5.3-43.5)	0.001*	6.5(2.6-26.8)
Current Alcohol use						
Yes	4	4	0.02*	6.2(1.4-26.6)	0.95	1.1(0.13-8.6)
No	20	123	Reference	Reference	Reference	Reference
Current Caffeine use						
Yes	19	94	0.59	0.8(0.26-2.2)		
No	5	33	Reference	Reference	Reference	Reference
Current Smoking cigarette						
Yes	5	7	0.02*	4.5(1.3-15.7)	0.04*	1.4(2.6-7.8)
No	19	120	Reference	Reference	Reference	Reference
Current Khat use						
Yes	13	57	0.4	0.69(0.29-1.65)		
No	11	70	Reference	Reference	Reference	Reference

*statistically significant

Table 3: Factors associated with antipsychotic induced TD among psychiatric patients on first-generation antipsychotics at JUSH, psychiatric clinic, Jimma, Ethiopia, 2016.

4. Discussion

This study investigated the prevalence and associated factors of Tardive dyskinesia (TD) among psychiatric patients on first-generation antipsychotics at Jimma University specialized hospital (JUSH) psychiatric clinic, Jimma, South west Ethiopia. The prevalence of TD in the current study was 14.6% with the range between 10.76% and 18.4%.

Our study result revealed that age, current smoking cigarettes and chlorpromazine equivalent dose range > 400 mg/day had statistically significant association with first-generation antipsychotic induced Tardive Dyskinesia (TD). In this study the prevalence of TD among psychiatric patients on first-generation antipsychotics is closer to the previous finding in Ethiopia which revealed that 11.9% of patients had TD [1]. On the other hand, the prevalence is greater than that found in Arab among psychotic patients treated with first-generation antipsychotic which shows the prevalence was 5.9% [9]. The present study result is also lower than the findings of studies done in Turkey and Indian [13, 14] in which the prevalence of first-generation antipsychotics induced TD was 22.5% and 26.4% respectively. This variation could be due to different possible reasons like characteristics of the participants. In the

current study we included patients with different diagnosis including schizophrenia, bipolar disorder and MDD with psychotic feature those were getting first-generation antipsychotic drugs. But studies like in Turkey and Indian did on only schizophrenic patients. The other point is also in the current study we include patients who took medications at least for six (6) months and above. But some the compared studies like study in Indian included participants who took medications for at least one (1) year and above.

Our study revealed that patients who had smoked cigarettes at least once during the last one month were 1.4 times more likely to develop TD as compared to patients who were not smoking cigarettes during the last one month (AOR 1.4, 95% CI: 2.6, 7.8) and this was similar with the result of previous studies which showed the prevalence of TD was significantly higher among smokers than non-smokers [15]. Out of the first-generation antipsychotics included in this study, Haloperidol, chlorpromazine, and thioridazine are metabolized by CYP enzymes except trifluoperazine which is metabolized by UDP-glucuronosyltransferase1A4. Even so, Smoking is associated with induction of CYP1A2, the study in Chinese revealed that there is no association between the C→A genetic polymorphism of CYP1A2 and TD [15]. Contrary to our study result, a study conducted in the eastern cape of south Africa showed that a history of smoking was not associated with an increased or decreased risk, although fewer subjects with TD were smokers and they had smoked for fewer years [16].

In our study patients who were in the age category > 45 year were more than four times more likely to develop TD as compared to patients who were found in the age category of <30 year (AOR 4.5, 95%CI: 9.7, 20.4) and it is in line with the results of many previous studies which revealed that the prevalence of TD is high among older patients [1, 9, 17-19]. This result indicates that whenever the professionals prescribe first-generation antipsychotics, they have to consider the patients' age and appropriately prescribe the medications as described in prescription guidelines. For instance, our study result showed that one of the associated factors for TD is getting relatively high dose of medications especially chlorpromazine. Therefore, the professionals could consider the patients' age and adjust the dose in order to prevent the TD.

The current study result shows that patients who had been getting chlorpromazine equivalent dose range > 400 mg/day were 6.5 times more likely to develop TD as compared to patients who had been getting chlorpromazine equivalent dose range 50-<100 mg/day (AOR 6.5, 95% CI: 2.6, 26.8) and this result was consistent with some previous study results [1, 16, 20]. This result forwards the message that the treating professionals (clinicians) have to put the patients on the effective lowest dose as much as possible. From our clinical experience, some times the treating clinicians will put the patients on unnecessary high dose which could expose the patients to different side effects including antipsychotics induced movement disorders.

The study has some limitations including: it was cross-sectional study and cannot show the cause-effect relationship between Tardive Dyskinesia and independent variables and also the independent factors included in this study were not comprehensive enough. Questions about some variables like current substance use (alcohol, Khat and cigarettes) are sensitive issue and social desirability bias is possible.

5. Conclusion

A considerable number of psychiatric patients were suffering from first-generation antipsychotics induced TD. Age, current smoking cigarettes and chlorpromazine equivalent dose were found to be significantly associated with first-generation antipsychotics induced TD. The study finding will provide information to form rational foundation for prevention and planning to bring change in contributing factors for developing TD. Also, the result could serve as base line information for further study. The study recommended that stakeholders should consider and appropriately address these statistically identified associated factors in order to reduce/prevent antipsychotics induced TD. Also, it is beneficial if professional care givers give attention for patients who are sensitive to antipsychotics like old age patients during treating them.

Competing Interests

The authors declare that they have no competing interests.

Authors' Contributions

TA contributed to the designing research, conduct and analysis of the research, drafting the manuscript, critically reviewed and approved the manuscript for publication. All authors read and approved the final manuscript. DT contributed to the design, conduct and analysis of the research and in the review of the manuscript. WT contributed to the designing research, conduct and analysis and in the review of the manuscript.

Acknowledgements

We would like to acknowledge Jimma University, institute of health for financial support of our study from data collection to analysis and finally we thank the study participants, data collectors, and supervisors.

References

1. Taye H, Awoke T, Ebrahim J. Antipsychotic medication induced movement disorders: The case of Amanuel Specialized Mental Hospital, Addis Ababa, Ethiopia.
2. Albayrak Y, Ekinçi O. Duloxetine-associated tardive dyskinesia resolved with fluvoxamine: a case report. *Journal of clinical psychopharmacology* 32 (2012): 723-724.
3. Carbon M, et al. Tardive dyskinesia risk with first-and second-generation antipsychotics in comparative randomized controlled trials: a meta-analysis. *World Psychiatry* 17 (2018): 330-340.
4. Margolese HC, et al. Tardive dyskinesia in the era of typical and atypical antipsychotics. Part 2: Incidence and management strategies in patients with schizophrenia. *The Canadian Journal of Psychiatry* 50 (2005): 703-714.
5. Cornett EM, et al. Medication-induced tardive dyskinesia: a review and update. *Ochsner Journal* 17 (2017): 162-174.
6. Lieberman JA, et al. Atypical and conventional antipsychotic drugs in treatment-naive first-episode schizophrenia: a 52-week randomized trial of clozapine vs chlorpromazine. *Neuropsychopharmacology* 28 (2003): 995.

7. Tarsy D, Baldessarini RJ. Epidemiology of tardive dyskinesia: is risk declining with modern antipsychotics? *Movement disorders: official journal of the Movement Disorder Society* 21 (2006): 589-598.
8. van Harten PN, et al. Incidence of tardive dyskinesia and tardive dystonia in African Caribbean patients on long-term antipsychotic treatment: the Curaçao extrapyramidal syndromes study V. *The Journal of clinical psychiatry* 67 (2006): 1920.
9. Osman A, et al. Neuroleptic-induced tardive dyskinesia among Arab psychotic patients/La dyskinesie tardive post-neuroleptique chez des patients arabes psychotiques. *Eastern Mediterranean Health Journal* 13 (2007): 625-633.
10. Lane RD, et al. Assessment of tardive dyskinesia using the Abnormal Involuntary Movement Scale. *Journal of Nervous and Mental Disease* (1985).
11. Schooler NR, Kane JM. Research diagnoses for tardive dyskinesia. *Archives of General Psychiatry* 39 (1982): 486-487.
12. Abdeta T, et al. Prevalence, withdrawal symptoms and associated factors of khat chewing among students at Jimma University in Ethiopia. *BMC psychiatry* 17 (2017): 142.
13. Nalbant A, et al. Tardive Dyskinesia in Long Term Hospitalized Patients with Schizophrenia/Uzun süreli ve yatirilarak izlenen sizofreni hastalarinda tardiv diskinezi. *Dusunen Adam* 29 (2016): 259.
14. Achalia RM, et al. Prevalence and risk factors associated with tardive dyskinesia among Indian patients with schizophrenia. *Asian journal of psychiatry* 9 (2014): 31-35.
15. Chong SA, Tan EC, Tan CH. Smoking and tardive dyskinesia: lack of involvement of the CYP1A2 gene. *Journal of Psychiatry and Neuroscience* 28 (2003): 185.
16. Patterson BD, Swingler D, Willows S. Prevalence of and risk factors for tardive dyskinesia in a Xhosa population in the Eastern Cape of South Africa. *Schizophrenia research* 76 (2005): 89-97.
17. Kane JM, Woerner M, Lieberman J. Tardive dyskinesia: prevalence, incidence, and risk factors. *Journal of clinical psychopharmacology* 8 (1988): 52S-56S.
18. Sadock BJ, Sadock VA. *Kaplan and Sadock's synopsis of psychiatry: Behavioral sciences/clinical psychiatry*. Lippincott Williams & Wilkins (2011).
19. Jeste DV. Tardive dyskinesia in older patients. *Journal of Clinical Psychiatry* 61 (2000): 27-32.
20. Jeste DV, et al. Incidence of tardive dyskinesia in early stages of low-dose treatment with typical neuroleptics in older patients. *American Journal of Psychiatry* 156 (1999): 309-311.

Citation: Tilahun Abdeta, Daniel Tolessa, Wondale Tsega. Prevalence and Associated Factors of Tardive Dyskinesia Among Psychiatric Patients on First-Generation Antipsychotics at Jimma University Specialized Hospital, Psychiatric Clinic, Ethiopia: Institution Based on A Cross-Sectional Study. *Journal of Psychiatry and Psychiatric Disorders* 3 (2019): 179-190.



This article is an open access article distributed under the terms and conditions of the [Creative Commons Attribution \(CC-BY\) license 4.0](https://creativecommons.org/licenses/by/4.0/)