

**STUDIES ON PHYSICO CHEMICAL CHARACTERISTICS OF CHIKKAMALAPPANAKERE  
TANK, SHIVAMOGGA, KARNATAKA, INDIA**

Mahesh Anand Goudar

Department of Chemistry, D.V.S.College of Arts and Science, Kuvempu University, Shivamogga- 577201,  
Karnataka, INDIA

E-mail - dvs.mahesh@gmail.com Mobile - 9482173962

**ABSTRACT:** An investigation was carried out in Chikkamalappanakere tank near Shivamogga town on physico-chemical characteristics during January to December 2012. The results of physico-chemical parameters were compared with the standard values prescribed by the Bureau of India Standards (BIS) and World Health Organization (WHO). The study revealed that, tank water is polluted as it possesses high BOD, CO<sub>2</sub>, phosphate and nitrogen. In the light of standard of water quality recommended by WHO, the tank water should not be used by human beings especially for drinking and cooking.

**Key Words:** Chikkamalappanakere tank, Physico-chemical parameters

**INTRODUCTION**

Water is precious and is the basic need for all living being on this planet. Ponds and tanks are important multi-usage components, they are sources of irrigation, fishery and other domestic purposes (Hocioglu and Dulger, 2011). Water is an essential component of the environment and it sustains life on the earth. Due to over expanding population and industrial settlements, the demand for fresh water is increasing day by day. In today's scenario, unplanned urbanization, rapid industrialization and indiscriminate use of artificial chemicals cause heavy and varied pollution in aquatic environments leading to deterioration of water quality and depletion of aquatic fauna. Even the stagnation of water in ponds and tanks causes severe problems in aquatic ecosystem and in damaging the water quality. Physico-chemical parameters play a vital role in determining the distributional pattern and quantitative abundance of organisms inhabiting a particular aquatic ecosystem (Santhoshkumar Singh *et al.*, 2009). Several investigators have studied the physico-chemical dynamics of varied lentic water bodies with the intent to assess the water quality (Begum *et al.*, 2010; Sayeswara *et al.*, 2010; Begum *et al.*, 2011; Purushothama *et al.*, 2011; Purushothama *et al.*, 2011a; Sayeswara *et al.*, 2011; Sayeswara *et al.*, 2011a; Sayeswara *et al.*, 2011b; Mahesh Anand Goudar & Sayeswara, 2011; Sayeswara *et al.*, 2012; Vasantha Naik *et al.*, 2013; Sayeswara, 2014; Mahesh Anand Goudar and Sayeswara, 2014).

**Study area**

Chikkamalappanakere tank is an annual water body situated at about 35 km away from Shivamogga, located between 15° 63'N latitude and 14° 4' E longitude. The tank receives water from the adjacent paddy fields, and rain is the main source of water. The river basin of the tank is Krishna river. The total area of Chikkamalappanakere tank is about 25 acres, of which water spreads over an area of 15 acres with an average depth of 8-10 feet. The water has undergone moderate changes in its physico-chemical properties due to ecological degradation, overflowing of water from adjacent paddy fields and other excessive human activities. The basis of selection of Chikkamalappanakere tank was that its water is used by a large population which receives adequate waste water and periodic flooding from plains. In the present investigation, an attempt has been made to assess the suitability of water for human consumption and domestic purposes.

## MATERIALS AND METHODS

Water was sampled on monthly basis, between 7 to 9 am from January to December 2012. This water samples were collected in good quality polythene bottles. Water temperature was recorded at the sampling site itself. Dissolved oxygen was fixed on the spot itself in BOD bottles. Various parameters like turbidity, total hardness, sulphate, free CO<sub>2</sub>, alkalinity, BOD, TDS, phosphate, nitrate and chloride were estimated as per the standard methods (APHA, 1998).

## RESULTS AND DISCUSSION

The results of monthly and seasonal variation of physico-chemical parameters of Chikkamalappanakere tank are given in Table 1 and depicted in figures 1-4.

**Temperature:** The water temperature is largely influenced by local climatic conditions. The seasonal water temperature ranged from 20.1 to 26.8°C. The minimum value was recorded in June and maximum in April. Turbidity is a measure of cloudiness of water. Turbidity in natural water arises due to the presence of suspended matter such as clay, silts, finely divided organic and inorganic matter, phytoplanktons and other microscopic organisms. The values of turbidity ranged from 9.3 to 59.1 NTU. The highest and the lowest values were recorded in July and March, respectively.

**pH:** pH values are slightly acidic to slightly alkaline and found within permissible limit of 6.5 to 8.5 as per the Bureau of Indian Standards (BIS, 1993). The minimum value was observed during January (6.3) and maximum during July (7.6). The pH is important since aquatic organisms are well adapted to specific pH range and do not withstand abrupt changes in it.

**Dissolved oxygen:** Dissolved oxygen is an important gaseous factor that determines the quality of water and intern regulates the distribution of aquatic organisms. In the present study the DO level fluctuated between 1.7 to 3.8 mg/L. The highest and the lowest values were recorded in August and April, respectively. The variations of DO depend on the primary production and respiration of aquatic organisms. The permissible standard of DO is above 5mg/L (Perk and Park, 1980).

**Biological Oxygen Demand:** BOD is the measure of degradable organic matter present in water. BOD and other microbial activities generally increase by the introduction of sewage (Hynes, 1971). In the present study BOD values ranged between 6.2 to 9.4 mg/L. The minimum value was noticed in the month of July while maximum in September. They were found above the permissible limit of 6.5mg/L (WHO, 1991).

**Carbon dioxide:** Free carbon dioxide values fluctuated between 10.7 to 28.6 mg/L. The highest and the lowest values were recorded in March and January, respectively. The variation of CO<sub>2</sub> was due to the absorption by plants for photosynthesis and activity of other living organisms.

**Alkalinity:** Alkalinity in the water samples is primarily a function of carbonate, bicarbonate and hydroxide content. In the present study total alkalinity ranged from 50.9mg/L (November) to 92.1mg/L (January). It is within permissible limit of 600mg/L (BIS, 1993). Surface alkalinity may result from the discharge of domestic wastes.

**Table 1. Physico-chemical characteristics of Chikkamalappanakere tank water.**

Parameters	Months:2012											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Temperature	23.2	22.8	23.8	26.8	20.2	20.1	22.3	21.7	23.3	22.1	22.8	20.1
Turbidity	11.2	10.2	10.1	107	28.4	78.4	60.9	33.9	33.9	21.6	17.4	11.2
pH	6.3	6.5	6.5	7.2	7.1	6.5	7.6	7.1	7.3	7.3	7.2	7.1
DO	2.1	2.2	1.9	1.7	2.4	2.8	3.2	3.8	3.1	2.6	2.4	2.7
BOD	7.2	7.1	6.8	7.4	8.3	7.1	6.2	7.2	9.4	8.2	7.7	7.9
CO <sub>2</sub>	10.7	15.7	27.4	20.1	21.5	19.2	16.1	22.5	17.3	18.7	21.7	28.6
Alkalinity	92.1	62.4	68.8	84.5	71.3	68.4	71.1	65.9	6.28	59.1	50.9	52.6
TDS	33.6	25.5	41.9	60.6	41.1	44.2	31.1	41.1	36.8	34.3	42.7	44.9
TH	91	81	91	101	124	83	80	69	89	71	86	91
Chloride	26.7	72.9	81.3	93.4	95.2	92.4	64.6	63.4	60.2	71.1	73.3	80.7
Phosphate	1.1	0.9	1.6	1.2	1.4	1.6	1.7	1.1	1.3	1.2	2.2	1.7
Nitrate	6.1	5.3	6.2	6.2	7.4	6.4	6.3	6.1	5.3	4.9	6.1	6.6
Sulphate	10.3	9.1	7.4	7.7	10.2	8.2	6.9	8.4	8.7	11.1	9.3	7.2

All values are expressed in mg/l except pH, temperature (°C) and turbidity (NTU)

**Total Dissolved Solids:** TDS values ranged from 34.3 to 10.6mg/L, the minimum was recorded in October and maximum in April. The minimum value may be due to the stagnant condition of the water body. The values are within permissible limits of 1500 mg/L (BIS, 1993).

**Total Hardness:** Total hardness of water is not a pollution parameter but indicates water quality mainly in terms of  $Ca^{2+}$  and  $Mg^{2+}$  contents. Total hardness values observed are 69 to 124 mg/L. The minimum value was recorded in October and maximum in May.




Figure 1. Physico-chemical characteristics of Chikkamalappanakere tank




Figure 2. . Physico-chemical characteristics of Chikkamalappanakere tank




Figure 3. Physico-chemical characteristics of Chikkamalappanakere tank




Figure 4. Physico-chemical characteristics of Chikkamalappanakere tank

**Chloride:** Chloride is an important anion found in variable amounts in water bodies. Chlorides increase the degree of eutrophication (Goel *et al.*, 1980). In the present study, chloride values fluctuated between 63.4 mg/L (September) to 95.2 mg/L (May). High chloride content indicates the deterioration of water quality usually linked with sewage load (Mini *et al.*, 2003). The most important sources of chlorides in the fresh water are the discharge of domestic sewage and farm drainage. The concentration of chlorides is thus the indicator of pollution.

**Phosphorus:** Phosphorus occurs in natural water as various types of phosphates. The most important sources of phosphates are the discharge of domestic sewage, detergents and agricultural runoff. Values of phosphates ranged from 0.9 to 2.2 mg/L with the minimum value in February and maximum in November.

**Nitrates:** Most of the unpolluted sources of water are deficient of nitrates because it exists only in few natural sources (Trivedi and Goel, 1984). In the present study, nitrate values ranged from 4.9 to 7.2 mg/L. The minimum value of nitrate was noticed in the month of October while maximum in May.

**Sulphate:** Sulphate is one of the major anions occurring in natural waters. It may enter natural waters through weathering of sulphate bearing deposits. The values fluctuated between 6.9 to 11.1 mg/L. The minimum value was recorded in July and maximum is January.

The water samples from Chikkamalappanakere tank was collected and analyzed for various physico-chemical parameters to study the extent of pollution. Planktonic composition was also studied. DO was very low and BOD, CO<sub>2</sub>, phosphate and nitrogen values were significantly higher than the permissible level for domestic consumption. In the light of standard of water quality recommended by WHO, the tank water should not be used by human beings especially for drinking and cooking. Precautionary measures should be taken before the water is consumed.

## ACKNOWLEDGEMENT

The authors express their gratitude to Dr. B.M. Hosur, Principal, D.V.S. College of Arts and Science, Shivamogga and for facilities and encouragement.

## REFERENCES

- APHA, (1998). Standard Methods for the Examination of Water and Waste Water, 20<sup>th</sup> ed. Public Health Association, Washington, D.C.
- BIS: 3025. (1993). Methods of sampling and Test (Physical and Chemical) for water and waste water, 1st Revision, 1-2.
- Goel, P. K., Gopal, B. and Trivedi, R.K. (1980). Impact of sewage on freshwater ecosystem, *Int. J. Ecol and Environ. Sci.*, 6: 97-116.
- Hocioglu, N. and Basaran Dulger. (2011). Monthly variation of some Physico-chemical and microbiological parameters in Biga Stream (Big, Canakkale, Turkey). *African J. Biotechnology*, 8(9): 1929-1937.
- Hynes, H. B. N. (1971). *The Biology of Polluted Water*. Uni. Toronto Press, Canada.

- Mahesh Anand Goudar and Sayeswara, H.A. (2011). Hydrochemistry of Bhudhigere tank near Shivamogga, Karnataka, India, *Current Biotica*, 5(1): 85-90.
- Mahesh Anand Goudar and Sayeswara, H.A. (2014). Studies on surface water quality evaluation of Gowrikere tank, Anandapura, Sagara, Shivamogga, Karnataka, India, *International Journal of Science and Nature*, 5(3): 567-571.
- Mini, I., Radhika, C. J. and Tunga Devi, T. (2003). Hydrobiological studies on a Lotic Ecosystem, Vamanapuram River, Thiruvananthapuram, Kerala, *Poll. Res.*, 22(4): 617-626.
- Nafeesa Begum, Narayana, J and Sayeswara, H.A. (2010). A seasonal study of Phytoplankton diversity and pollution indicators of Bathi pond near Davangere city, Karnataka (India). *Environmental Conservation Journal*, 11(3): 75-80.
- Nafeesa Begum, Sayeswara, H.A. and Naik, K.L. (2011). Seasonal variations of Phytoplankton diversity in Bethur pond near Davangere, Karnataka, *Environment & Ecology*, 29(3): 1355-1359.
- Perk, J. E. and Park, K. E. (1980). *A Text book of preventives and social medicine*, 8<sup>th</sup> edition, Messer Banrsidas Bhanot, Jabalpur.
- Purushothama, R, Mahesh Anand Goudar and Sayeswara, H.A. (2011). Seasonal Phytoplankton diversity and diversity in two lentic water bodies of Sagara, Karnataka, India, *International Journal of Chemical Sciences*, 9(3): 1373-1390.
- Purushothama, R, Sayeswara, H.A. and Mahesh Anand Goudar (2011a). Dynamics of zooplankton diversity in relation to water quality of Heggere tank, Kanale, Sagara, Karnataka, India, *Environmental Conservation Journal*, 12(1&2): 29-34.
- Santhokumar Singh, A. Dakua, D. and Biswas, S. P. (2009). Physico-chemical parameters and fish enumeration of Majjan Beel (Wetland) of Upper Assam, *Geobios*, 36: 184-188.
- Sayeswara, H.A, Naik, K.L, Nafeesa Begum and Ashashree, H.M. (2011). Potability of water in relation to some Physico-chemical parameter of Mudugodu pond, Chikkamagalur, Karnataka, India, *Environment & Ecology*, 29(1): 140-142.
- Sayeswara, H.A, Mahesh Anand Goudar and Manjunatha, R (2011). Water quality evaluation of Phytoplankton diversity Hosahalli tank, Shivamogga, Karnataka (India). *International Journal of Chemical Sciences*, 9(2): 805-815.
- Sayeswara, H.A., Ravikumar patil, H.S. and Mahesh Anand Goudar (2010a). Studies of Physico-chemical parameters of Purla pond water of Shivamogga, Karnataka (India). *International Journal of Chemical Sciences*, 8(1): 582-588.
- Sayeswara, H.A, Vasantha Naik, T, Ravikumar Patil, H.S. and Mahesh Anand Goudar. (2012). Study on Habitat ecology and Phytoplankton diversity of Nagathibelagalu tank in industrial town Bhadravathi, Karnataka, India, *Nature, Environment and Pollution Technology*, 11(3): 419-423.
- Sayeswara, H.A.(2014). Phytoplankton species diversity in Chikkamalappanakere tank, Shivamogga, Karnataka, India, *International Journal of Life sciences, Biotechnology and Pharma research*, 3(4): 98-102.
- Shinde, S.E., Pathan, T.S., Raut, K.S. and Sonawane, D.L. (2001). Studies on Physico-chemical parameters and correlation coefficient of Harsool-savangi dam, Aurangabad, India, *Middle-East Journal of Scientific Research*, 8(3): 544-554
- Trivedi, R.K., Goel, P. K. and Trishal, C. L. (1995). *Practical methods in Ecology and Environmental Science*, Environmental Publications, Karad (India).
- Vasanth Naik, T, Puttaiah, E.T, Ravikumar Patil, H.S, Sayeswara, H.A. and Nafeesa Begum. (2013). Phytoplankton community composition in Gowrikere tank, Anandapura, Sagara, Shivamogga, Karnataka, India, *Science, Technology and Arts Journal*, 2(2): 70-74.
- WHO, (1991). *International Standards for drinking water*, Geneva.



ISSN : 0976-4550

# INTERNATIONAL JOURNAL OF APPLIED BIOLOGY AND PHARMACEUTICAL TECHNOLOGY



Email : [ijabpt@gmail.com](mailto:ijabpt@gmail.com)

Website: [www.ijabpt.com](http://www.ijabpt.com)