

**SOME BIOCHEMICAL CHANGES IN SERUM OF MALE ALBINO RATS TREATED WITH
AQUEOUS LEAF EXTRACT OF *PHYLLANTHUS AMARUS***

¹Chinyere, Godwin Chinyere ¹Uhegbu, Friday Obinwa, ¹Emmanuel Iroha AKUBUGWO, ²Ositadinma Chinyere UGBOGU and ¹Amadike Eziuche UGBOGU*

Department of ¹Biochemistry and ²Microbiology, Abia State University Uturu, Nigeria

Corresponding Author e-mail: amasryal@yahoo.com. Tel: +2348053574147

ABSTRACT : Biochemical Changes in serum of male albino rats treated with aqueous leaf extract of *Phyllanthus annus* were examined. 50mg/kg and 100mg/kg of the extracts were administered orally and once daily to group I and group II of test animals respectively for 28 days. The third group which served as control received distilled water only. On the 29th day, the rats which had been fasted overnight were dissected under chloroform anaesthesia and blood was collected directly from their heart. The blood was allowed to clot and centrifuged to obtain serum which was kept in a refrigerator at - 20°C until its usage for the analysis of the following parameters; alanine and aspartate transaminases, alkaline and acid phosphatases, haemoglobin concentration, high and low density lipoproteins. The result indicates significant ($P < 0.05$) increases in AST, ALT and ACP levels in tests groups compared to the control animals. There were non-significant elevation in the concentration of ALP, HDL, LDL and haemoglobin. These results suggest that *P. amarus* may be toxic if abused but if properly administered can be use for its medicinal values.

Keywords: *Phyllanthus annus*, liver enzymes, lipoproteins, rats.

INTRODUCTION

The plant *Phyllanthus amarus* belongs to the family Euphorbiaceae, which is a family otherwise, called the spurge family. It is a large family of about 300 genera and 600 species (Heywood, 1978). Most members are trees, shrubs and few are herbs; of rainforest and xerophylactic habitats (Burkill, 1994). Although plants of this family have important economic uses as foodstuffs, medicinal and industrials, particularly as sources of rubber and timber, nonetheless most members of this family are poisonous (Adedapo, 2002).

Phyllanthus amarus presently is considered as one of the best herbs for treating disorders such as hepatitis B, diabetes mellitus, skin ulcer (Oluwafemi and Debiri, 2008; Joseph and Raj, 2011). The plant has diuretic and purgative action and is also known to have a remedy for inflammation of the respiratory tract and for asthma as it has a special reputation for causing bronchial relaxation (Odetola and Akejenu, 2000; Raphael and Kuttan, 2003; Kuttan, 2004, Adeneye et al, 2006). The main constituents of the plant include lignins, alkaloids, flavonoids, phenols and terpenes (Gotto, 1998; Joseph and Raj, 2011).

This study was aimed to investigate the effect of *Phyllanthus amarus* on some liver enzymes and lipoproteins of albino rats so as to adequately explore its toxicity at therapeutic doses.

MATERIALS AND METHODS**Plant collection and extraction**

Fresh leaves of *P. amarus* were collected within Abia State University and were taken to the department of Biochemistry. The plant was identified by a taxonomist. The leaves were sorted to remove the dead ones washed without squeezing to remove debris and dust particles. Large quantities of the leaves were collected and sun-dried for 5 days. The dried leaves were milled to get a coarse powder used for the extraction. Exactly 50g of the powder were macerated in a percolator with 250ml of distilled water. The mixture was allowed to stand for 24 hours after which it was filtered. The filtrate was then placed in an oven to evaporate and the solid residue (5.0g) referred to as extract. 5.0g of the residue was added to 250ml (2.5% solution w/v) of water and administered to rats orally.

Experimental Animals

Twenty four white male albino rats (*Rattus norvegicus*) weighing between 105 to 120g were obtained from small animal breeding unit of the Department of Biochemistry Abia State University Uturu. The rats had free access to water and standard palletized feed and they were kept in the care of experienced animal technicians. They were randomly divided into three (3) groups with each group containing eight (8) rats.

Administration of Extract

Rats in group I were fed 100 mg/kg of the extract and Rats in group II were fed 50 mg/kg of the extract orally once daily for 28 days to each groups while the control received distilled water only instead of the extract. Animals were thereafter exposed to feed and water *ad libitum*.

Collection of Blood: On the 29th day, blood was collected directly from the heart of the overnight fasted rats under chloroform anaesthesia. The collected blood was allowed to clot and centrifuged to obtain the serum. The serum was kept in a refrigerator at 0-20°C until further use.

Estimation of Biomolecules

Appropriate commercial Kits (Randox Laboratories, U.K) were used to determine the concentrations of alanine and aspartate transaminases (ALT and AST), alkaline and acid phosphatases (ALP and ACP), and cholesterol and haemoglobin.

ALT and AST were estimated using the method of Reitman and Frankel, 1957 based on the principle that ALT involves the monitoring of the concentration of pyruvate hydrazone formed with 2,4-dinitrophenyl hydrazine while AST involved monitoring the concentration of oxaloacetate hydrazone formed 2,4-dinitrophenyl hydrazine.

Alkaline phosphatase activity of the serum was determined as described by Bessey *et al.*, (1946), based on the principle that serum alkaline phosphatase hydrolyzes a colourless substrate of phenolphthalein that result in phosphoric acid and phenolphthalein at alkaline pH values. Pink coloured product is measured calorimetrically at 550nm.

Total cholesterol: This was carried out by the enzymatic colorimetric chod-PAP method.

High density lipoprotein (HDL)-Cholesterol: HDL separated from chylomicrons. Very low density lipoprotein (VLDL) and Low density lipoproteins (LDL) are separated by the addition of a phosphotungstic and magnesium chloride (precipitating reagent) to the serum (Burstein *et al.*, 1970). After centrifugation, the cholesterol content was determined by the enzymatic colorimetric method.

Haemoglobin was determined as described by Dacie and Lewis, 1990.

Statistical Analysis

All analysis was performed in triplicates. Results were expressed in mean±S.D. Statistical significance was established using Analysis of variance (ANOVA). Means were Separated according to Duncan's multiple range analysis ($P < 0.05$).

RESULTS

Table 1 shows the body weight gain and haemoglobin concentration in rats of different groups during entire experimental period. The weight of the group I and group II were not differ significantly ($P < 0.05$) as compared to the control. whereas haemoglobin concentration was found to be maximum in group I animals as compared to other group. The result in Table 2 shows a Significant ($P < 0.05$) increased activities of AST and ALP were observed in Group I and II animals as compared to the control animals. ALT were less than 20.00 IU/L for all the test and control animals and activity was significantly high in treated groups.

ACP has the highest value (16.16) in group II animals compared to (12.18) in Group I and in control animals (9.11). HDL-C for group I, group II and control animals were 26.90, 29.10 and 28.00 respectively and animals of group II are higher as compared to control and group I. The LDL level in group II (33.10 ± 6.60 Mg/L) was higher as compared to control (32.00 ± 0.10 Mg/L) and group I (33.40 ± 0.70 Mg/L) animals.

Table 1. EFFECTS OF *PHYTHANTHUS ANNUS* ON BODY WEIGHT AND HAEMOGLOBIN CONCENTRATION OF MALE ABINO RATS.

Parameters	Group I (administered) 1.5ml of the extract	Group 2 administered 1.0ml of the extract	Control
Body weight (g)	108.00 ± 2.00^a	110.80 ± 3.10^a	111.50 ± 2.50^a
Haemoglobin concentration (Mg/dL)	34.60 ± 0.30^b	31.20 ± 0.10^a	29.80 ± 0.40^a

Figures are mean ± S.D figures bearing different alphabets on the row differ significantly ($P < 0.05$) according to Duncan's multiple range analysis N = 8

Table 2 : EFFECTS OF *PHYTHANTHUS ANNUS* ON SOME SERUM BIOCHEMICAL PARAMETERS OF MALE ABINO RATS.

Parameters	Group I (administered) 1.5ml of the extract	Group 2 administered 10ml of the extract	Control
AST (IU/L)	46.30±3.00 ^c	33.70±2.40 ^b	29.6±2.20 ^a
ALT (IU/L)	18.80±2.00 ^b	18.40±1.00 ^b	16.80±2.10 ^a
ALP (IU/L)	46.06±1.24 ^b	46.16±3.40 ^b	43.68±2.00 ^a
ACP (IU/L)	14.18±0.46 ^a	16.16±0.79 ^a	12.11±0.45 ^a
HDL-C(Mg/L)	30.90±0.90 ^a	29.10±0.6 ^b	28.00±0.01 ^a
LDL (Mg/L)	33.40±0.70 ^b	33.10±6.60 ^{ab}	32.00±0.10 ^a

Figures are mean ± S.D figures bearing different alphabets on the row differ significantly (P<0.05) according to Duncan's multiple range analysis, N = 8.

DISCUSSION

All the animals gained weight and the weight gained were not significant (P<0.05) compared to the control.

Haemoglobin functions physiologically only in transport of oxygen from the living to the peripheral tissues and also in transportation of carbon dioxide from the tissues to the lungs (Tietz, 1999). Significant increase (P<0.05) in haemoglobin concentration observed group 1 (Table I) which occurred as the concentration of the extract increases may result in the lowering of oxidative stress (Gupta *et al.*; 2002).

AST is an enzyme that is present in high quantities in the cytoplasm and mitochondria of liver, heart, skeletal muscle, kidney and brain (Benjamin, 1978; Ringler and Dabrich, 1999). Elevation in the activity of AST can be associated with cell necrosis of many tissues. For example, pathology involving the skeletal or cardiac muscle and hepatic parenchyma allows for the leakage of large, amounts of this enzyme into the blood (Kaneko, 1980; Duncan *et al.*, 1994). The elevation in AST produced by this plant is an indication of tissue necrosis.

ALT is cytosolic enzyme is useful in measuring hepatic necrosis, especially in small animals (Bush, 1991). Since it is one of the specific liver enzymes its elevated level in this study may indicate hepatic damage by this plant.

ALP is reported to be present in large number of cells but only in a few cells in the activity sufficient to be of clinical importance. It is found in liver cells and is associated with osteoblastic activity in bone (saini and saini, 1978), it means that these quantities may contain little toxic component compared to the control and may not cause injury to several tissues of the body.

ACP found throughout the body but primarily in the prostate gland (Muller *et al.*, 2000) and aqueous extract of *Phyllanthus amarus* has no effect on the activity of ACP on these doses.

HDL major function is to remove unesterified cholesterol where it may have accumulated in cell membranes and plasma lipoprotein and transport it to the liver where it can be degraded and utilized for among other things the synthesis of the bile acids (Guyton, 1981) while LDL transports cholesterol to tissues where it may be needed for membrane structure or conversion into various metabolites such as steroid hormones (Gurr and Harwood, 1991). There were slight increases in HDL and LDL tests compared to the control animals but these increases were not significant (P<0.05). This indicates that *P. amarus* has no negative impact on HDL and LDL levels in the test animals.

CONCLUSION

Aqueous extract of *Phyllanthus annus* significant (P<0.05) increase in AST and ALT while other serum biochemical parameters are non-significantly alter at lower doses in male albino rats. The results of the present study suggest that higher doses of the plant extract may produces damaging effects in different tissues.

REFERENCES

- Adedapo, A.A (2002). Toxicological effect of some plants in the family Euphorbiaceae on rats. Ph.D thesis University of Ibadan.
- Adeneye, A.A., Amole, O.O. and Adeneye, A.K. (2006). Hypoglycemic and hypocholesterolemic activities of the aqueous leaf and seed extracts of *Phyllanthus amarus* in mice. *Fitoterapia*, 77: 511-514.
- Benjamin M.N (1978). Outline of veterinary clinical pathology. University Press. 10WA, USA. Pp 229-232.
- Burkill, H.M (1994): The useful plants of West Tropical Africa. Royal Botanical Gardens, Ken.

- Burstein M., Scholnick H.R., Morfin R. (1970). Rapid method for isolation of lipoproteins from human serum by precipitation with polyanions. *J Lipid Res*, 11 , 583- 595
- Bush, B.M (1991): Interpretation of laboratory results for small animal clinicians. Blackwell scientific Publications Oxford.
- Cornelius, C.E (1989): Serum enzyme activities and other markers for detecting hepatic necrosis, cholestasis or carcinoma in clinical Biochemistry of Domestic Animals Academy press Toronto, PP. 381-386.
- Dacie, J.V. and Lewis, S.M. (1990). Practical Hematology. Churchill Livingstone New York 34.
- Duncan, J.R; Prasse, k.W ; and Mahaffey, E.A (1994). Veterinary laboratory medicine (clinical Pathology). Iowa state University Press.
- Gotto A.M (1988). Lipoprotein Metabolism and the ethiology of hyperlipidemia, Hospital practice 23:1-4.
- Gupta, S.K; Prakash J and Srivastava, S (2002). Validation of traditional claim of Tulsi, *Ocimum Sanctum* linn as a medicinal plant. *Indian J. Exp. Bio*, 40:765-773
- Gurr, M.I and Harwood, J.L (1991). Lipid Biochemistry. An introduction fourth edition Chapman and Hall. Pp 200-210
- Guyton, A.C (1981). Textbook on medical Physiology. 6 edition. W.B saunders company Philadelphia, Pp 463-474.
- Heywood, V. H (1978). Flowering plants of the world. Oxford University Press.
- Joseph, B. and Raj, S.J. (2011). An Overview: Pharmacognostic Properties of *Phyllanthus amarus* Linn.. *Inter J. Pharmacol.*, 7: 40-45.
- Kaneko, J.J (1980). In Kaneko, J.J (Ed) Clinical Biochemistry of Domestic Animals. American Press. Orlands, Fla Pp 21-24.
- Kuttan, KB (2004). Prospective effect of an extract of *Phyllanthus amarus* against radiation-induced damage mice. *J. Radiat. Res*, (45(1):133-139.
- Maller PB; Taboada J, Hosgood G; Partington B.P vansteenhout J.L; Taylor H. W (2000). Effects of longterm Phenobarbital treatment on the liver in dogs. *J. vet. Inter. Med*. 14:165-171.
- Odetola, A.A and Akojenu, S.M (2000). Antidiarrhoeal and gartrointertinal potentials of the aqueous extract of *Phyllanthus amarus*. *J. Ethnopharmacol.*, 81(1): 17-22.
- Oluwafemi, F. and F. Debiri, (2008). Antimicrobial effect of *Phyllantus amarus* and *Parquetina nigrescens* on *Salmonella typhi*. *Afr. J. Biomed. Res.*, 11: 215-219
- Raphael K.R and Kuttan R. (2003). Inhibition of experimental gastric lession and inflammation by *Phyllanthus amarus* extract. *J. Ethnopharmacol.*, 87(2): 193-197.
- Reitman, S. and Frankel, S.L. (1957). Determination of ALT and AST concentrations in serum. *Am. J. Chem Path.*, 28:56
- Ringler, D.H and Dabich, L. (1979). Haematology and clinical Biochemistry. In the laboratory Rat. Vol 1 edited by II.J. Baker; J.R lindsey, S.H. Weisbroth. Academic Press. London. Pp 105-118.
- Saini P, K and Saini, B.K (1978). origin of Serum alkaline phosphatase in the Dog. *Am. J. vet. Res*. 39:1510-1513.
- Tietz, N.W (1999). Clinical guide to laboratory test, 3rd W.B. Saunders company. Philadelphia Pp 578-510.