

## Research Article

# Seperation of bioactive compounds from Haemolymph of scarab beetle *Scarabaeus sacer* (Coleoptera: Scarabaeidae) by GC-MS and determination of its antimicrobial activity

Nancy Taha Mohamed<sup>1</sup>, Doaa Hassan Abdelsalam<sup>1</sup>, Ahmed Salem El-Ebiarie<sup>1</sup>, Mahmoud Elaasser<sup>2</sup>

<sup>1</sup>Zoology & Entomology Department, Faculty of Science, Helwan University

<sup>2</sup>Regional Centre of Mycology and Biotechnology, Al-Azhar University

**\*Corresponding author:** Doaa Hassan Abdelsalam, Zoology & Entomology Department, Faculty of Science, Helwan University, Egypt.

**Received:** 20 November 2021; **Accepted:** 02 December 2021; **Published:** 07 December 2021

**Citation:** Nancy Taha Mohamed, Doaa Hassan Abdelsalam, Ahmed Salem El-Ebiarie, Mahmoud Elaasser. Seperation of bioactive compounds from Haemolymph of scarab beetle *Scarabaeus sacer* (Coleoptera: Scarabaeidae) by GC-MS and determination of its antimicrobial activity. International Journal of Applied Biology and Pharmaceutical Technology 12 (2021): 461-480.

### Abstract

This study aimed to investigate the bioactive compounds in the haemolymph of scarab beetle *Scarabaeus sacer* by using the Gas chromatography–mass spectrometry (GS-MS) analysis. The identification of the bioactive compounds is based on peak area, retention time, molecular formula and molecular weight. There are 129 compounds are detected in the haemolymph of scarab beetle and 43 of them were reported to have a bioactivity. The most analyzed bioactive compounds are alcohols, steroids, fatty acids and terpenoids. The current study also test the antimicrobial activity of scarab beetle haemolymph against gram-negative bacteria (*Escherichia coli*, *Enterobacter cloacae*, gram-positive bacteria (*Staphylococcus aureus*, *Bacillus subtilis*), and fungi (*Aspergillus fumigatus* and *Candida albicans*). The haemolymph has highest antibacterial activity against gram negative bacteria *Enterobacter cloacae*,

*Escherichia coli* respectively and against gram-positive bacteria *Bacillus subtilis*, *Staphylococcus aureus* respectively. No antifungal activity has been detected.

**Keywords:** Haemolymph, GS-MS, gram negative bacteria, gram positive bacteria, antimicrobial activity

## 1. Introduction

Coleoptera is one of the largest order of insects with about 370,000 insect species described worldwide. The family Scarabaeidae encompasses over 30,000 species of beetles worldwide; they are often called scarabs. Dung beetles are a major insect group (Coleoptera: Scarabaeidae) distributed globally except Antarctica with a high number of diversity comprising approximately 6,200 species and nearly 267 genera [1]. These species are coprophagous in nature which live freely in soil and mostly feed on both wet and dry dung materials of herbivorous mammals. The undigested excreta of mammals are utilized as food and nesting material throughout their life cycle, hence, they possess many ecologically beneficial functions.

The dung beetles play a vital role in nutrient recycling by decaying organic matter and developing soil aeration [2] thereby, reducing the greenhouse gas fluxes [3]. It also improves plant growth and grain production [4]. *Scarabeus sacer* is considered a species of genus *Scarabaeus*, occurs in coastal dunes and marshes around the Mediterranean Basin. It can be found across North Africa, Southern Europe and parts of Asia (Afghanistan, Corsica, Cyprus, France, Iran, Israel, Italy, Morocco, Sardinia, Sicily, Sudan and Syria). *Scarabaeus sacer* is a species of dung beetle belonging to the family Scarabaeidae [5].

Insects are known for their ability to resist infection. They protect themselves against bacterial infection by secreting a battery of antimicrobial peptides (AMP) into the hemolymph. Hemolymph, also known as the insect blood, is a clear fluid, with or without yellow or greenish pigmentation. It constitutes 16-40% of the body weight of certain insects. The volume and component of hemolymph are vary in different types of insects and their developmental stages. It spends much of its time flowing freely within body cavities where it makes direct contact with all internal tissues and organs. Therefore the circulation would help to transport the AMP to its target site [6]. In insects, AMPs / polypeptides are manufactured mainly in a fat body (similar to mammalian liver) and are released into hemolymph where they play a vital role in innate immune systems and host defense mechanisms, and having a broad spectrum of activity against both gram + ve and gram -ve bacteria and against fungi [7].

However, misuse of antibiotics intake has caused many problems, such as the appearance of antibiotic-resistant bacteria, weakening of disease resistance in livestock, and ecosystem pollution [8]. Insects exhibit innate immune systems that produce potent AMPs to protect them from pathogen invasion, and these AMPs are viewed as strong natural antibiotic applicants [9]. The insect innate immune system is categorized into cellular and humoral immunity. Cellular immunity involves the phagocytosis of bacteria, fungi, and protozoa, and nodule formation and encapsulation, while humoral immunity involves the secretions of proteins and peptides produced in fat and blood cells to hemolymph in response to infection [10].

AMPs secreted by the humoral immune response are classified according to their structure and amino acid sequence into cecropins, defensins, proline-rich peptides, glycine-rich peptides, and lysozymes and are found in various insect orders including Coleoptera, Diptera, Hymenoptera, and Lepidoptera [11]. Melittin is familiar AMP contained in bee venom and its antimicrobial activity was observed greatly in methicillin-resistant *Staphylococcus aureus* (MRSA) and Gram-positive and Gram-negative bacteria [12]. AMPs are small molecules that vary in size, ranging from 10 to 100 amino acid residues and are produced by all living organisms. The rich diversity of insects makes them rich sources of AMPs. The black soldier fly *Hermetia illucens* L. (Diptera: Stratiomyidae), particularly, able to live in hostile environments rich in microbial colonies, making it one of the most promising sources of AMPs [13].

Nowadays, the using of GC-MS technique is important in analyzing and separation the compounds found in plants extracts [14] and also the haemolymph of arthropods. This will be a new method for discovering a future drug to be used in traditional medicine system. In this article, the haemolymph of scarab beetle was analyzed by GC-MS and resulting in many bioactive compounds.

## 2. Material and Methodology

### 2.1 Collection of Insects

Scarab beetle was collected from Baltim in the Kafr El Sheikh Governorate, in the north coast of Egypt.

### 2.2 Withdrawing of Hemolymph

Scarab beetle body surface cleaned with 70% alcohol. Then, in order to collect haemolymph, hind pair legs were cut from coxa, and haemolymph fluid was extracted with a capillary tube placed into micro tubes containing EDTA. Haemolymph was centrifuged at  $10000 \times g$  for 10 minutes and the supernatant was collected for the antimicrobial testing and stored in 4°C.

### 2.3 Gas Chromatography–Mass Spectrometry (GC-MS) Analysis

Haemolymph was collected and 30 milligrams were homogenized in 1ml methanol centrifuged at 4500 rpm for 10 minutes, the supernatant was taken to GC-MS. The chemical composition of samples was performed using Trace GC1310-ISQ mass spectrometer (Thermo Scientific, Austin, TX, USA) with a direct capillary column TG–5MS (30 m x 0.25 mm x 0.25 µm film thickness). The column oven temperature was initially held at 50 C and then increased by 5°C /min to 230°C for 2 min. increased to the final temperature 290°C by 30°C /min and hold for 2 min. The injector and MS transfer line temperatures were kept at 250, 260°C respectively; Helium was used as a carrier gas at a constant flow rate of 1 ml/min. The solvent delay was 3 min and diluted samples of 1 µl were injected automatically using Auto-sampler AS1300 coupled with GC in the split mode. EI mass spectra were collected at 70 eV ionization voltages over the range of m/z 40–1000 in full scan mode. The ion source temperature was set at 200 °C. The components were identified by comparison of their retention times and mass spectra with those of the WILEY 09 and NIST 11 mass spectral databases.

## 2.4 Antimicrobial Activity Assay

The antimicrobial activity was investigated in the haemolymph against microorganisms. All microbial strains were provided from the culture collection of the Regional Center for Mycology and Biotechnology (RCMB), Al-Azhar University, Cairo, Egypt. The antimicrobial profile was tested against two Gram-positive bacterial species (*Bacillus subtilis*, *Staphylococcus aureus*), two Gram-negative bacterial species (*Escherichia coli*, *Enterobacter cloacae*) and two fungi (*Aspergillus fumigatus* and *Candida albicans*) using a modified well diffusion method. Briefly, 100 µl of the test bacteria/or fungi were grown in 10 mL of fresh media until they reached a count of approximately  $10^8$  cells/ml for bacteria or  $10^5$  cells/mL for fungi [15]. One hundred µl of microbial suspension was spread onto agar plates corresponding to the broth in which they were maintained and tested for susceptibility by well diffusion method on Mueller-Hinton and Sabaroud agar (Clinical and Laboratory Standards Institute, 2012.) One hundred µL of each sample (at 10 mg/ml) was added to each well (10 mm diameter holes cut in the agar gel). The plates were incubated for 24-48 h at 37 °C (for bacteria and yeast) and for 48 h at 28 °C (for filamentous fungi). After incubation, the microorganism's growth was observed.

The resulting inhibition zone diameters were measured in millimeters and used as a criterion for antimicrobial activity. If an organism is placed on the agar, it will not grow in the area around the well if it is susceptible to the chemical. This area of no growth around the disc is known as a "Zone of inhibition" or "Clear zone". The size of the clear zone is proportional to the inhibitory action of the compound under investigation. Solvent controls (DMSO) were included in every experiment as negative controls. DMSO was used for dissolving the tested compounds and showed no inhibition zones, confirming that it has no influence on growth of the tested microorganisms. Gentamycin and ketoconazole (Sigma Aldrich, USA) were used as standard antibacterial and antifungal drugs at 30 and 50ug/ml, respectively.

### 2.4.1 MIC Determination

The tested extract was screened in vitro for their antibacterial and antifungal activities at a different concentration to determine the lowest concentration inhibiting the growth of the organism that recorded as the MIC [15].

## 3. Results and Discussion

In the present article, the separation of compounds in the haemolymph of scarab beetle by using GC-MS analysis gas separation technique resulting in 43 bioactive compounds as shown in table (1). The identification of the bioactive compounds is based on peak area, retention time, molecular formula and molecular weight.

In this study, the antibacterial effectiveness of the haemolymph against gram -V bacteria (*Escherichia coli*, *Enterobacter cloacae*, against gram +V bacteria (*Staphylococcus aureus*, *Bacillus subtilis*) as well as antifungal activity (*Aspergillus fumigatus*, *Candida albicans*) is investigated. Antibacterial activity against *Bacillus subtilis* was first, followed by *Enterobacter cloacae*, *Escherichia coli* and *Staphylococcus aureus* respectively. No antifungal activity has been investigated yet (Table 2). The minimum inhibitory concentration (MIC) is measured for the tested

sample and it was 2500 µg/ml against *E. coli*, 1250 µg/ml against *Enterobacter cloacae*, *Bacillus subtilis* and 10000 µg/ml against *Staphylococcus aureus* respectively (Table 3).

**Table 1:** Bioactive compounds in haemolymph of scarab beetle separating by GC-MS

No.	Compound Name	Molecular Formula	Chemical nature	MWT	Area %	RT	Bioactivity
1	Trichloromethane	CHCl <sub>3</sub>	Trihalomethane	118	3.57	4.06	Anti-virus, anti-cancer, anti-mutagenic, anti-allergic and anti-ulcer [16]
2	Tert-Hexadecanethiol	C <sub>16</sub> H <sub>34</sub> S	Thio-Alcohol	258	0.11	5.16	Enzyme activators [17]
3	Ethanol, 2 octadecyloxy-	C <sub>20</sub> H <sub>42</sub> O <sub>2</sub>	Alcohol	314	0.11	5.16	Antimicrobial activity [18]
4	1-Heptatriacotanol	C <sub>37</sub> H <sub>76</sub> O	Alcohol	536	0.48	39.87	Anticancer, antineoplastic and anti-HIV [19]
5	1,4-Benzenediol, 2-(1,1-dimethylethyl)-5-(2-propenyl)-	C <sub>13</sub> H <sub>18</sub> O <sub>2</sub>	Alcohol	206	2.66	25.14	Anticancer, Antioxidant activity and pesticides [20]
6	E,E,Z-1,3,12-Nonadecatriene-5,14-diol	C <sub>19</sub> H <sub>34</sub> O <sub>2</sub>	Alcohol	294	0.40	39.20	Antimicrobial activity [21]
7	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C <sub>20</sub> H <sub>40</sub> O	Terpene alcohol	296	6.43	30.04	Antimicrobial [22] Precursor of synthetic forms of vitamin E and vitamin K1 [23]

8	Dotriacontane	C32H66	Saturated hydrocarbons	450	0.11	5.16	Antimicrobial, antifungal, anti-inflammatory, cytotoxic activity [24]
9	Isochiapin B	C19H22O6	Terpenoids	346	0.32	10.83	Anti-insect Antitumor agent [25]
10	14- $\alpha$ -H-Pregna	C21H36	Steroids	288	0.32	10.83	defense chemical and Diabetic retinopathy prevention [26]
11	Cycloheptasiloxane, tetradecamethyl-	C14H42O7Si7	Organo-Silicone compound	518	1.69	15.25	Antimicrobial, Antiseptic, Hair Conditioning Agent, Skin-Conditioning Agent-Emollient and Solvent [27]
12	4H-1-Benzopyran-4 one, 2-(3,4-dimethoxyphenyl)-3, 5-dihydroxy-7-methoxy-	C18H16O7	Ketonic comound	344	0.38	17.48	Antioxidant, antimicrobial, cancer enzyme inhibitors in pharmaceutical, cosmetics, and food industries [28]
13	Cyclooctasiloxane, hexadecamethyl-	C16H48O8Si8	Alkanes	592	2.84	19.68	Antimicrobial [29]
14	Cyclododecasiloxane, tetracosamethyl	C24H72O12Si12	Alkanes	888	2.84	19.68	Hepatoprotective, antispasmodic

							antirheumatic [29]
15	9,12-Octadecadienoic acid (Z,Z)-, 2,3-bis [(trimethylsilyl)oxy]propyl ester	C27H54O4Si2	Linoleic acid ester	498	0.74	21.58	Anti-inflammation [30]
16	4H-1-Benzopyran-4-one, 2-(3,4-dimethoxyphenyl)-3, 5-dihydroxy-7-methoxy-	C18H16O7	Phenolic compound	344	0.28	25.36	Antioxidant, antimicrobial, cancer enzyme inhibitors in pharmaceutical, cosmetics, and food industries [31]
17	9,12,15-Octadecatrienoic acid ,2,3-bis[(Trimethylsilyl)oxy]propyl ester, (Z,Z,Z)-	C27H52O4Si2	Alpha linoleic acid	496	0.28	25.36	anti-oxidant, anti-diabetic, and anti-inflammatory [15]
18	Eicosamethylcyclodocasiloxane	C20H60O10Si10	Organoheterosilane	740	2.59	27.05	Prevent degenerative diseases [32]
19	Silicone oil	N/A	polysiloxane compounds	0	2.59	27.05	Toxic for bed-bug(insecticide) [33]
20	1H-Purin-6-amine,[(2-fluorophenyl)methyl]	C12H10FN5	Amino compounds	243	2.59	27.05	Anti-oxidant [32]
21	Neophytadiene	C20H38	Alkenes	278	6.43	30.04	analgesic, antipyretic, anti-inflammatory, antimicrobial,

							and antioxidant compound [34]
22	2,6,10-Trimethyl, 14-ethylene-14-pentadecene	C <sub>20</sub> H <sub>38</sub>	Alkenes	278	6.43	30.04	Antiproliferative activity [21]
23	Phytol Isomer	C <sub>20</sub> H <sub>40</sub> O	Diterpene	296	0.29	30.93	antimicrobial, antioxidant, and anticancer activities [28]
24	7-Methyl-Z-tetradecen-1-ol acetate	C <sub>17</sub> H <sub>32</sub> O <sub>2</sub>	Acetate ester	268	0.64	31.36	Anticancer, anti-inflammatory, hepatoprotective [35]
25	2-Dodecen-1-yl(-)succinic anhydride	C <sub>16</sub> H <sub>26</sub> O <sub>3</sub>	Succinic Acid anhydride	266	0.13	33.83	Antineoplastic agents, Antioxidants, Antimicrobial [36]
26	Hexadecanoic acid, methyl ester	C <sub>17</sub> H <sub>34</sub> O	Palmitic acid ester	270	2.09	34.34	Anti-oxidant, decrease blood cholesterol, anti-inflammatory [37]
27	Pentadecanoic acid, 14-methyl-, methyl ester	C <sub>17</sub> H <sub>34</sub> O	Ester	270	2.09	34.34	Antimicrobial, antifungal [38]
28	Dasycarpidan-1-methanol, (Ester)	C <sub>20</sub> H <sub>26</sub> N <sub>2</sub> O <sub>2</sub>	Ester	326	0.74	21.58	Antimicrobial [28]
29	Dibutyl phthalate	C <sub>16</sub> H <sub>22</sub> O <sub>4</sub>	Benzoic acid ester	278	0.98	38.03	Use in cosmetics [39] ectoparasitici



							de [40]
30	Phthalic acid, butyl undecyl ester	C23H36O4	Ester	376	0.98	38.03	Antimicrobial Antibacterial & Anti-inflammatory [41]
31	Phthalic acid, butyl tetradecyl ester	C26H42O4	Ester	418	0.98	38.03	Antimicrobial activity [42]
32	11,14-Eicosadienoic acid, methyl ester	C21H38O2	Ester	322	0.40	39.20	Anti-inflammatory, anti-oxidant, anti-arthritic, anti-coronary. [43]
33	1,2-Benzenedicarboxylic acid, butyl decyl ester	C22H34O4	Ester	362	5.04	39.35	Antimicrobial activity [44]
34	Diisooctyl phthalate	C24H38O4	Phthalic acid derivative (Ester)	390	1.80	50.09	Antimicrobial activity [45]
35	9,12,15-Octadecatrienoic acid, 2,3-dihydroxypropyl ester, (Z,Z,Z)-	C21H36O4	Linoleic acid ester	352	0.87	52.61	Analgesic, Antipyretic, Anticonvulsant, Antiseptic [46]
36	9,12,15-Octadecatrienoic acid, 2-phenyl-1,3-dioxan-5-yl ester	C28H40O4	Phenolic ester	440	0.46	58.55	Antimicrobial & anti-inflammatory [47]
37	Cholest-5-en-3-ol, 24-propylidene-, (3 $\alpha$ )-	C30H50O	Fatty acid	426	6.09	63.51	Antibacterial activity [48]
38	Cis-13-Eicosenoic acid	C20H38O2	Fatty acid	310	0.08	33.78	Anti-inflammatory activity [49]
39	Propionic acid, 3-(1-hydroxy-2-isopropyl-	C13H20O3	Fatty acid	224	1.72	37.54	Anti-angiogenic

	5 methylcyclohexyl)-						activity against solid tumor growth [50]
40	2-Nonadecanone 2,4-dinitrophenylhydrazine	C <sub>25</sub> H <sub>42</sub> N <sub>4</sub> O <sub>4</sub>	Nitrogen compound	462	0.36	50.66	Antimicrobial activity [51]
41	Ethyl iso-allocholate	C <sub>26</sub> H <sub>44</sub> O <sub>5</sub>	Steroid derivative	436	0.48	39.87	Antimicrobial, Antioxidant, anti-inflammatory & anti-arthritis antiasthmatic [52]
42	1,2-15,16-Diepoxyhexadecane	C <sub>16</sub> H <sub>30</sub> O <sub>2</sub>	Epoxide	254	0.40	39.20	Antitumor, anti-inflammatory [53]
43	Milbemycin B, 6,28-anhydro-15-chloro-25-isopropyl-13-dehydro-5-O-demethyl-4-methyl	C <sub>33</sub> H <sub>47</sub> ClO <sub>7</sub>	-----	590	0.33	58.77	Anti MRSA [54]

The GC-MS analysis on the haemolymph of *Scarabeus sacer* revealed presence of some bioactive compounds such as alcohols, terpenoids, ketones, phenolic compounds, alkanes, alkenes, amino-compounds, Fatty acids and steroids. Alcohols were discovered to have antimicrobial activity [55]. Isochiapin B, 3, 7, 11, 15-Tetramethyl-2-hexadecen-1-ol and Phytol Isomer are terpenoids present in haemolymph of *S. sacer*. Terpenoid compounds (Phorbol, Isochiapin B, stigmasterol acetate, and  $\beta$ -sitosterol) were detected in essential oil of *Achillea fragmentissima* that well known for their biological activities as anti-insect and anti-tumor agents [23]. Terpenes are bioactive compounds detected in *Ulva fasciata*, *U. lactuca* and *Corallina mediterranea* seaweeds extract and steroids were detected in the extracts of *U. fasciata*, and *Amphiroa anceps* seaweeds [56]. Most of these compounds exhibit biological activities such as anticancer, antiviral, antioxidant, and anti-inflammatories [57].

Phytol isomer is a diterpenes identified in the haemolymph of adult *S. sacer*. phytol is a bioactive compound that has a potent anticancer activity [58]. It also serve as a chemical attractant for parasitoids, according to research on these species: *Lucilia sericata* [55], *Leptinotarsa decemlineata* [65]. Alkanes can help distinguish organisms by acting as a chemical signal (Lockey 1988). Alkanes were also marked in the surface lipids of *Liposcelis bostrychophila*, *Cryptolestes ferrugineus* [60] and *Laelius utilis* (Howard, 1992). Dotriacontane is saturated hydrocarbons present in haemolymph of scarab beetle and reported to has antimicrobial, antifungal, anti-inflammatory and cytotoxic activity

[22]. Also the hydrocarbons used to distinguish between the male and female of *Sarcophaga* species [59]. Alkanes were also marked in the surface lipids of *Liposcelis bostrychophila*, *Cryptolestes ferrugineus* [60] and *Laelius utilis* [61]. Larvae of potato beetle contain hydrocarbons of high molecular weight, particularly tetrapentacontane (C<sub>54</sub>H<sub>110</sub>), pentapentacontane (C<sub>55</sub>H<sub>112</sub>) and heptapentacontane (C<sub>57</sub>H<sub>116</sub>) [62].

Pentadecanoic acid, 9, 12, 15-octadecatrienoic acid, hexadecanoic acid methyl ester were detected in the haemolymph of adult *S. sacer*. Pentadecanoic acid and 9, 12, 15-octadecatrienoic acid reported to have anti-inflammatory, antimicrobial, antioxidants, and antiproliferative activity [28]. Hexadecanoic acid methyl ester is also known as palmitic acid ester and efficiently used as an antioxidant, pesticide, anti-androgenic, nematocide, flavoring agent, hypocholesterolemic, and lubricant [63]. Also hexadecanoic acid was the major fatty acid in *Sargassum granuliferum* seaweed which prevents the biofilm forming bacteria [64].

Esters with even longer C<sub>22</sub>, C<sub>24</sub>, C<sub>42</sub> and C<sub>46</sub> carbon chains were determined in *Aleurothitius timberlakei* [65]. In the haemolymph of scarab beetle, fatty acids such as – Cis-13-eicosenoic acid, Cholest-5-en-3-ol, 24-propylidene-(3á), and Ppropionic acid, 3-(1-hydroxy-2-isopropyl-5 methylcyclohexyl) were detected. In case of *Leptinotarsa decemlineata* (Coleoptera: Chrysomelidae), the presence of C<sub>6</sub>, C<sub>9</sub>, C<sub>10</sub>, C<sub>12</sub>, C<sub>14</sub>, C<sub>16</sub> and C<sub>18</sub> fatty acids were found. Fatty acids C<sub>18</sub> and C<sub>20</sub> are found also in *Bombyx mori* and *Blatella germanica* [66]. Fatty acids C<sub>18</sub>:1, C<sub>18</sub>:2 and C<sub>18</sub>:3 were estimated in the larvae of *Drosophila melanogaster*, *Musca domestica* and *Galleria mellonella* [67]. Fatty acids C<sub>16</sub> – C<sub>18</sub> have also been determined in the surface lipids of *Cryptolestes ferrugineus* and *Liposcelis bostrychophila* insects [60]. Cholest-5-en-3-ol, 24-propylidene have been detected in the methyl extract of *Sargassum crassifolium* [67 and [68] separated Cholest-5-en-3-ol, 24-propylidene from the red alga *Grateloupia turuturu*.

In the adult scarab beetle haemolymph, there is a ketonic compounds such as 4H-1-Benzopyran-4 one, 2-(3, 4-dimethoxyphenyl)-3,5-dihydroxy-7-methoxy that act as Antioxidant, antimicrobial, cancer enzyme inhibitors in pharmaceutical, cosmetics, and food industries [26]. Also ketones are separated from *Tessaratomia papillosa* [69]. The relationships of these analyzed compounds in insects play vital roles as they can be transmitters of information and signals [70] and also serve as pheromones [71]. Ethyl iso-allocholate is a steroid derivative compounds detected in the haemolymph of adult scarab beetle and in the black fruit of *Pistacia lentiscus*, this steroid compound has antimicrobial, anti-inflammatory, anticancer, antiasthma and diuretic activities [72]

Silicone oil, milbemycin B, 6, 28-anhydro-15-chloro-25-isopropyl-13-dehydro-5-O-demethyl-4-met-, and Dibutyl phthalate were detected in the haemolymph of adult scarab beetle. [32], reported that silicone oil in the bed bug is cytotoxic and has an insecticidal activity that can kill insects by physical mean that affecting on tracheal system causing asphyxiation of insects. Anti MRSA activity was reported for milbemycin B, 6, 28-anhydro-15-chloro-25-isopropyl-13-dehydro-5-O-demethyl-4-met- [54] and Dibutyl phthalate act as ectoparasiticide [43].

**Table 2:** Antimicrobial activity (as a mean zone of inhibition) of the haemolymph of adult scarab beetle

Sample	Heamolymph	Control
Tested microorganisms		
<u>FUNGI</u>		<i>Ketoconazole</i>
<i>Aspergillus fumigatus</i>	NA	17
<i>Candida albicans</i>	NA	20
<u>Gram Positive Bacteria:</u>		<i>Gentamycin</i>
<i>Staphylococcus aureus</i>	8	24
<i>Bacillus subtilis</i> RCMB 015 (1)	15	26
<u>Gram Negatvie Bacteria:</u>		<i>Gentamycin</i>
<i>Escherichia coli</i>	10	30
<i>Enterobacter cloacae</i>	14	27

\*NA: No activity

**Table 3:** The antimicrobial activity as Minimum Inhibitory concentration (MIC) in µg/ml of the tested microorganisms. The test was done using the diffusion agar technique

Sample	Heamolymph	Control
Tested microorganisms		
<u>Gram Positive Bacteria:</u>		<i>Gentamycin</i>
<i>Staphylococcus aureus</i>	10000	3.9
<i>Bacillus subtitle's</i>	1250	1.95
<u>Gram Negatvie Bacteria:</u>		<i>Gentamycin</i>
<i>Escherichia coli</i>	2500	1.95
<i>Enterobacter cloacae</i>	1250	3.9

\*NA: No activity

The present works approved that the haemolymph of *S. sacer* possesses antibacterial activity against gram –negative bacteria (*Escherichia coli*, *Enterobacter cloacae*) and against gram-positive bacteria (*Staphylococcus aureus*, *Bacillus subtilis*). No antifungal activity had been investigated against *Aspergillus fumigatus* and *Candida albicans*. There are many works hassling to our work; the methanol extract of oriental hornet *Vespa orientalis* and *Zophobas mori* (Coleoptera:Tenebrionidae) larva show antibacterial activity against *E. coli* and no antifungal activity [73]. Contrary to our results, the whole body extract of housefly maggots show no activity against *E. coli* and exhibit antifungal activity [74]. While [75] documented that the extract of the housefly maggots have higher activity against Gram- positive bacteria than Gram negative bacteria and had not antifungal activity yet.

Most of insect extracts show antibacterial activity against Gram-positive and Gram-negative bacteria, the silk worm *Bombyx mori* [76], the European bumble bee, *Bombus pascuorum* [77] and *Tenebrio molitor* larvae [78]. On the other hand, some other insects revealed activity only against Gram-positive bacteria as *Aedes aegypti* [79], *Chironomus plumosus* [80] and *Anopheles gambiae* [81].

Synthetic antibiotics and antimicrobials have contributed to public health and stimulated the growth of livestock. Conversely, overuse and abuse of antibiotics and antimicrobial drugs may cause drug-resistant bacteria, which threaten public and livestock health. Several studies reported that insects manufacture antimicrobial peptides (AMPs) which act as a natural antibiotic [82]. Insects not only perform different roles in the environment, but also host a variety of community of microorganisms. The complicated cellular and humoral mechanisms include the innate immune system of an insect [83]. The cellular mechanism is rely on phagocytosis process which is activated by enzymes and invading microorganisms then encapsulated by the hemolymph. Moreover, The humoral response is represented in the production of broad-spectrum antimicrobial peptides (AMPs), reactive oxygen or nitrogen intermediates, and complex enzymatic cascades that help to regulate hemolymph coagulation or melanization [84]. The presence of microorganisms invading insects causes the fat body to rapidly synthesize AMPs, which are then released into the hemolymph [85]. Previous research shows that each insect species produces a distinct antimicrobial peptide that acts against specific microorganisms [86]. On the other hand, in order to enhance the insect's defense system against other pathogens, some of the peptides are expressed simultaneously, encouraging synergism [87]. As such, AMPs have a specific modes of action, such as altering the electrochemical gradient at the membrane, producing reactive oxygen/nitrogen species (ROS/RNS) that cause cell death, inhibiting protein synthesis, and permeabilizing the cell membrane [88]. AMPs have pharmacological properties such as low molecular weight, high water solubility, broad-spectrum antimicrobial activity, and low levels of cytotoxicity [89]. [90] reported that the antibacterial activity in salivary secretions of *Polistes dominulus* larvae inhibits growth of Gram positive *Bacillus subtilis* and Gram negative *E. coli*. There are number of studies that have tested ability of the insect extracts against pathogenic bacteria, especially antimicrobial peptides extracted from various insects maggots [91], dung beetles [92], Red Palm Weevil [93], pupae of the giant silk moths [94].

Contrary to our study on the haemolymph of adult *S. sacer*, the non-induced hemolymph of dung beetle, *Onthophagus taurus* did not show inhibitory activity against any of the bacterial strains and fungus. It does not mean that peptides are absent but it may be present in smaller quantity so that no visible action in in-vitro studies is detected [95]. But the immune induced hemolymph exhibits activity against all tested bacteria and no activity against fungus. Therefore, the peptide is active against prokaryotes and doesn't affect the fungus which is a eukaryote. Many studies on insect species assert that bacteria injected into the haemocoel stimulate the synthesis of number of peptides and proteins which are active singly or in concert against the invaders and are secreted into the hemolymph [96].

#### 4. Conclusion

On conclusion, antimicrobial activity of haemolymph of adult *S. sacer* may be due to presence of the previous bioactive compounds which separated by the GC-MS technique. Future studies are necessary to purify the compounds with antimicrobial activity and investigate their antitumor effect against different cell lines.

#### Acknowledgement

Great thanks and appreciation for the Mycology Center, Al-Azhar University for their cooperation.

#### References

1. Ahmed MH Ali, Abdur Rauf, Emad M Abdallah. Insects as producers of antimicrobial polypeptides: A short review. GSC Biol. Pharm. Sci 12 (2020): 102–107.
2. Al-Gara’Awi NI, Abu-Serag NA, Alee Shaheed KA, Bahadly ZK. Al, 2019. Analysis of bioactive phytochemical compound of (*Cyperus alternifolius* L.) by using gas chromatography-mass spectrometry. IOP Conf. Ser. Mater. Sci. Eng (2019): 571.
3. Al Bratty M, Makeen HA, Alhazmi HA, Alhazmi HA, Syame SM, Syame SM, Abdalla AN, Abdalla AN, Homeida HE, Sultana S, Ahsan W, Khalid A, Khalid A. Phytochemical, Cytotoxic, and Antimicrobial Evaluation of the Fruits of Miswak Plant, *Salvadora persica* L. J. Chem (2020).
4. Albergoni VEPOC. esponse to heavy metals in organisms-I. Excretion and accumulation of physiological and non physiological metals in *Euglena gracilis*. Comp Biochem Physiol C Comp Pharmacol 67 (1980): 121–127.
5. Albratty M, Bajawi AAM, Marei TMH, Shamsheer Alam M, Alhazmi HA, Najmi A, ur Rehman Z, Sivagurunathan Moni S. Spectral analysis and antibacterial potential of bioactive principles of *Sargassum crassifolium* J. Agardh from Red sea of Jazan origin. Saudi J. Biol. Sci 28 (2021): 5745–5753.
6. Ali HAM, Imad HH, Salah AI. Analysis of bioactive chemical components of two medicinal plants (*Coriandrum sativum* and *Melia azedarach*) leaves using gas chromatography-mass spectrometry (GC-MS). African J. Biotechnol 14 (2015): 2812–2830.
7. Ali Shafaghat. Phytochemical and antimicrobial activities of *Lavandula officinalis* leaves and stems against some pathogenic microorganisms. J. Med. Plants Res 6 (2012): 455–460.
8. Ardenne M. von Osske G, Schreiber K, Steinfelder KR Tümmler. No Title. J. Insect Physiol 11 (1965): 1365.
9. Bakar K, Mohamad H, Latip J, Tan HS, Heng GM. Fatty acids compositions of *Sargassum granuliferum* and *Dictyota dichotoma* and their anti-fouling activities. J. Sustain. Sci. Manag 12 (2017): 8–16.
10. Barlow J. S., 1964. Fatty acids in some insect and spider fats. Can. J. Biochem. 42, 1365–1374.
11. Bekele D, Tekie H, Asfaw Z, Petros B. Bioactive Chemical Constituents from the Leaf of *Oreosyce africana* Hook.f (Cucurbitaceae) with Mosquitocidal Activities against Adult *Anopheles arabiensis*, the Principal Malaria Vector in Ethiopia. J. Fertil. Pestic 7 (2016): 1–8.
12. Beschi DA, Appavoo MR, Wilsy JI. GC-MS analysis, collected from Kavalkinaru area, Tirunelveli District, Tamil Nadu, India. Eur. J. Mol. & Clin. Med 7 (2021): 4287–4292.

13. Budayatin Waluyo J, Wahyuni D, Dafik. Antibacterial effects of *Pheretima javanica* extract and bioactive chemical analysis using Gas Chromatography Mass Spectrum. *J. Phys. Conf Ser* (2021) : 1751.
14. Chernysh S, Gordya N, Suborova T. Insect antimicrobial peptide complexes prevent resistance development in bacteria. *PLoS One* 10 (2015).
15. Chinnasamy PS, Parimala S, Kandhasamy M. Phytochemical Evaluation of Seed and Fruit Pulp EXTRACTS OF *PASSIFLORA FOETIDA* L P 7 (2018): 1924–1932.
16. Daffodil D Almeida, Mohan VR, Uthayakumari FK. GC-MS determination of bioactive compounds of *Curculigo orchioides* gaertn. *Sci. Res. Report 2* (2012): 198–201.
17. Devakumar J, Keerthana V, Sudha SS. Identification of bioactive compounds by gas chromatography-mass spectrometry analysis of *Syzygium jambos* (L.) collected from Western Ghats region Coimbatore, Tamil Nadu. *Asian J. Pharm. Clin. Res* 10 (2017): 364–369.
18. Durak D, Kalender Y. Fine structure and chemical analysis of the metathoracic scent gland of *Eurygaster maura* (Linnaeus, 1758) (Heteroptera: Scutelleridae). *Folia Biol. (Praha)* 55 (2007): 133–141.
19. Elsharkawy E. Anti-inflammatory activity and chemical compositions of essential oil of *Achillea fragmentissima*. *Natl. J. Physiol. Pharm. Pharmacol* 6 (2016): 258–262.
20. Erwan Plouguerné, Kikuchi H, Oshima Y, Deslandes E, Stiger-Pouvreau V. Isolation of Cholest-5-en-3-ol formate from the red alga *Grateloupia turuturu* Yamada and its chemotaxonomic significance. *Biochem. Syst. Ecol* 34 (2006): 714–717.
21. Eva de Rijke, Out P, Niessen WMA, Ariese F, Gooijer C, Brinkman UAT. Analytical separation and detection methods for flavonoids. *J. Chromatogr* 11 (2006): 31–63.
22. Gillespie JP, MR K TT. Biological mediators of insect immunity. *Annu Rev Entomol* 42 (1997): 611–643.
23. Gołębowski M, Dawgul M, Kamysz W, Boguś MI, Wieloch W, Włóka E, Paszkiewicz M, Przybysz E, Stepnowski P. Antimicrobial activity of alcohols from *Musca domestica*. *J. Exp. Biol* 215 (2012): 3419–3428.
24. Guo G, Wu J, Fu P, Qin R, Zhang Y, Song Z. Isolation and purification of antibacterial peptides from the larvae secretion of housefly and the characteristics. *Chinese J. Parasitol. Parasit Dis* 25 (2007): 87–92.
25. Hadi MY, Mohammed GJ, Hameed IH. Analysis of bioactive chemical compounds of *Nigella sativa* using gas chromatography-mass spectrometry. *J. Pharmacogn. Phyther* 8 (2016): 8–24.
26. Hameed IH, Altameme HJ, Idan SA. *Artemisia annua*: Biochemical products analysis of methanolic aerial parts extract and anti-microbial capacity *Res J. Pharm Biol Chem Sci* 7 (2016): 1843–1868.
27. Hameed IH, Hussein HJ, Kareem MA, Hamad NS. Identification of five newly described bioactive chemical compounds in Methanolic extract of *Mentha viridis* by using gas chromatography – mass spectrometry (GC-MS). *J. Pharmacogn. Phyther* 7 (2015): 107–125.
28. Harris ED. Symposium : Regulation of Antioxidant Enzymes Copper as a Cofactor and Regulator of Copper, Zinc Superoxide Dismutase 12 (1992): 636–640.
29. HASSAN MI, MOHAMED AF, AMER MS, KOTB M. HAMMAD, MAHBOUB MT. Antimicrobial Activity of Three Insect Species, Crude Extracts Againsts Certain Microbial Agents. *Al-Azhar Bull Sci* 26 (2015): 19–24.

30. Hema R, Kumaravel S, Alagusundaram Agardh K. GC - MS determination of bioactive components of *Gracilaria dura*. *J. Am. Sci* 5 (2015): 100–105.
31. Hoffmann JA, Reichhart JM. *Drosophila* innate immunity: An evolutionary perspective. *Nat. Immunol* 3 (2002): 121–126.
32. Hou L, Le YS, Zhai P, Guowei Le. Antibacterial activity and in vitro antitumor activity of the extract of the larvae of the housefly, *Musca domestica*. *J. Ethnopharmacol* 111 (2007): 227–231.
33. Howard RW. Comparative analysis of cuticular hydrocarbons from the ectoparasitoids *Cephalonomia waterstoni* and *Laelius utilis* (Hymenoptera: Bethyridae) and their respective hosts, *Cryptolestes ferrugineus* (Coleoptera: Cucujidae) and *Trogoderma variabile* (Coleoptera: Ann. Entomol. Soc Am 85 (1992): 317–325.
34. Howard RW, Lord JC. Cuticular lipids of the booklouse, *Liposcelis bostrychophila*: Hydrocarbons, aldehydes, fatty acids, and fatty acid amides. *J. Chem. Ecol.* 29 (2003): 615–627.
35. Huang D, Irwin G. Computational intelligence: international conference on intelligent computing. Springer-Verlag (2006): 16–19.
36. Hussein HJ, Hameed IH, Hadi MY. Using gas chromatography-mass spectrometry (GC-MS) technique for analysis of bioactive compounds of methanolic leaves extract of *Lepidium sativum*. *Res. J. Pharm. Technol* 10 (2017): 3981–3989.
37. Hussein HM, Hameed IH, Ibraheem OA. Antimicrobial activity and spectral chemical analysis of methanolic leaves extract of *Adiantum capillus-veneris* using GC-MS and FT-IR spectroscopy. *Int. J. Pharmacogn. Phytochem. Res* 8 (2016): 369–385.
38. Ibrahim H, Eldehna W, Abdel-Aziz H, Elaasser M, Abdel-Aziz M. Improvement of antibacterial activity of some sulfa drugs through linkage to certain phthalazin-1(2H)-one scaffolds. *Eur. J. Med Chem* 85 (2014): 480.
39. Januszani B, Stączek S, Zdybicka -Barabas A, Bądziul D, Jakubowicz-Gil J, Langner E, Rzeski W, Cytryńska M. The effect of *Galleria mellonella* polypeptides on human brain glioblastoma multiforme cell line – a preliminary study / Badania wstępne wpływu polipeptydów hemolimfy *Galleria mellonella* na komórki ludzkiego glejaka wielopostaciowego. *Ann. UMCS Biol* 67 (2013).
40. Jatin RR, Sonawani P. Determination of Bioactive Components of *Cynodon dactylon* by GC-MS Analysis & its In Vitro Antimicrobial Activity © Sakun Publishing House (SPH): IJPLS. *Int. J. Pharm. LIFE Sci. Int* 7 (2016): 4880–4885.
41. Jiang L, Wang W, He Q, Wu Y, Lu Z, Sun J, Liu Z, Shao Y, Wang A. Oleic acid induces apoptosis and autophagy in the treatment of Tongue Squamous cell carcinomas. *Sci Rep* 7 (2017): 1–11.
42. Kadhim MJ, Al-Rubaye AF, Hameed IH. Determination of Bioactive Compounds of Methanolic Extract of *Vitis vinifera* Using GC-MS. *Int. J. Toxicol. Pharmacol Res* 9 (2017): 113–126.
43. Kala SC, Ammani K. GC-MS analysis of biologically active compounds in *Canthium parviflorum* Lam. leaf and callus extracts *Ijergg* 10 (2017): 1039–1058.
44. Kalsy M, Tonk M, Hardt M, Dobrindt U, Zdybicka-Barabas A, Cytryńska M, Vilcinskas A, Mukherjee K. The insect antimicrobial peptide cecropin A disrupts uropathogenic *Escherichia coli* biofilms. *npj Biofilms*



- Microbiomes 6 (2020).
45. Kanost MR, Jiang H, Yu XQ. Innate immune responses of a lepidopteran insect, *Manduca sexta*. *Immunol. Rev* 198 (2004): 97–105.
  46. Karthikeyan AVP, Sudan I. Gc-Ms Profile of in Vivo and in Vitro Shoots of *Cleome gynandra* L. *Int. J. Pharm. Pharm. Sci* 9 (2017): 21.
  47. Koyama M, Iwata R, Yamane A, Katase T, Ueda S. Nutrient intake in the third instar larvae of *Anomala cuprea* and *Protaetia orientalis submarmorea* (Coleoptera: Scarabaeidae) from a mixture of cow dung and wood chips: Results from stable isotope analyses of nitrogen and carbon. *Appl. Entomol. Zool* 38 (2003): 305–311.
  48. Kurata S. Intra- and extracellular recognition of pathogens and activation of innate immunity. *Yakugaku Zasshi* 126 (2006): 1213–1218.
  49. Lauth X, Nesin A, Briand J, Roussee J, Hetru C. Isolation, characterization and chemical synthesis of a new insect defensin from *Chironomus plumosus* (Diptera). *Insect Biochem Mol Biol* 28 (1998): 1059–1066.
  50. Lee KH, Hong SY, Oh JE, Kwon M, Yoon JH, Lee J, Lee BL. Defensin of *Tenebrio molitor* 105 (1998): 99–105.
  51. Lei J, Sun LC, Huang S, Zhu C, Li P, He J, Mackey V, Coy DH, He QY. The antimicrobial peptides and their potential clinical applications. *Am. J. Transl. Res* 11 (2019): 3919–3931.
  52. Long G. No Title *The British Museum: Egyptian Antiquities, Volume 2 Library of entertaining knowledge The British Museum: Egyptian Antiquities, George Long. Charles Knight, New York (1836).*
  53. Looft T, Johnson TA, Allen HK, Bayles DO, Alt DP, Stedtfeld RD, Sul WJ, Stedtfeld TM, Chai B, Cole JR, Hashsham SA, Tiedje JM, Stanton TB. In-feed antibiotic effects on the swine intestinal microbiome. *Proc. Natl. Acad. Sci USA* 109 (2012): 1691–1696.
  54. Lowenberger C, Bulet P, Charlet M, Hetru C, Hodgeman B, Christensen BM, JAH. Insect immunity: isolation of three novel inducible antibacterial defensins from the vector mosquito, *Aedes aegypti*. *Insect Biochem Mol Biol* 25 (1995): 867–873.
  55. Manning P, Slade EM, Beynon SA, Lewis OT. Functionally rich dung beetle assemblages are required to provide multiple ecosystem services. *Agric. Ecosyst. Environ* 218 (2016): 87–94.
  56. Mary APF, Giri S. *World Journal of Pharmaceutical Research GC-MS ANALYSIS OF BIOACTIVE COMPOUNDS OF 7* (2018): 1045–1056.
  57. Mary Ann Liebert. Final Report on the Safety Assessment of Dibutyl Phthalate, Dimethyl Phthalate, and Diethyl Phthalate. *Journal of Am Coll Toxicol* 4 (1985): 267–303.
  58. Meylaers K, Clynen E, Daloz D, Deloof A, Schoofs L. Identification of 1-lysophosphatidylethanolamine (C(16:1)) as an antimicrobial compound in the housefly, *Musca domestica*. *Insect Biochem. Mol Biol* 34 (2004): 43–49.
  59. Michael Zasloff. Antimicrobial peptides of multicellular organisms *Nature* 415 (2002): 389–395.
  60. Mofeed J, Deyab M, Sabry AE-N, Ward F. In Vitro Anticancer Activity of Five Marine Seaweeds Extract From Egypt Against Human Breast and Colon Cancer Cell Lines. *Res Sq* 1 (2021): 1–15.
  61. Mohtar JA, Yusof F, Ali NMH. Screening of novel acidified solvents for maximal antimicrobial peptide extraction from *Zophobas morio fabricius*. *Adv Environ Biol* 8 (2014): 803–809.

62. Moore HE, Hall MJR, Drijfhout FP, Cody RB, Whitmore D. Cuticular hydrocarbons for identifying Sarcophagidae (Diptera). *Sci. Rep* 11 (2021): 1–11.
63. Moretta A, Salvia R, Scieuzo C, Di Somma A, Vogel H, Pucci P, Sgambato A, Wolff M, Falabella P. A bioinformatic study of antimicrobial peptides identified in the Black Soldier Fly (BSF) *Hermetia illucens* (Diptera: Stratiomyidae). *Sci. Rep* 10 (2020): 1–15.
64. Muthu lakshmi A, Jothibai Margret R, Mohan VR. GC-MS analysis of bioactive components of *Feronia elephantum* Correa (Rutaceae). *J. Appl. Pharm. Sci* 2 (2012): 69–74.
65. Nelson DR, Walker GP, Buukner JS, Fatland CL. Composition of the wax particles and surface wax of adult whiteflies: *Aleuroplatus coronata*, *Aleurothrixus floccosus*, *Aleurotithius timberlakei*, *Dialeurodes citiri*, *Dialeurodes citrifolii*, and *Parabemisia myricae*. *Comp. Biochem. Physiol* 117B (1997): 241–251.
66. Noguez JH, Conner ES, Zhou Y, Ciche TA, Ragains JR, Butcher RA. *NIH Public Access* 7 (2013): 961–966.
67. Pashaei F, Bevalian P, Akbari R, Pooshang Bagheri K. Single dose eradication of extensively drug resistant *Acinetobacter* spp. In a mouse model of burn infection by melittin antimicrobial peptide. *Microb. Pathog* 127 (2019): 60–69.
68. Patil HV, Kumar BS. Isolation and partial purification of antimicrobial peptides/ proteins from dung beetle, *Onthophagus Taurus* immune hemolymph. *J. Microbiol. Biotechnol Food Sci* 2 (2013): 2414–2418.
69. Ponmathi Sujatha A, Michael Evanjaline R, Muthukumarasamy S, Mohan V. Determination of bioactive components of *barleria courtallica* nees (Acanthaceae) by gas chromatography–mass spectrometry analysis. *Asian J. Pharm. Clin. Res* (2017).
70. Prabhu V, Devi V, Priya MK. Gc-Ms Analysis of Bioactive Compounds Present in the Petroleum Ether , Chloroform and Methanol Extract of *Ixora Coccinea* ' S Flower and in-Vitro Cytotoxic Activity of. *Int. J. Res. Anal. Rev* 5 (2018): 801–807.
71. Rahnamaeian M, Cytryńska M, Zdybicka-Barabas A, Dobszlaff K, Wiesner J, Twyman RM, Zuchner T, Sadd BM, Regoes RR, Schmid-Hempel P, Vilcinskis A. Insect antimicrobial peptides show potentiating functional interactions against Gram-negative bacteria. *Proc. R. Soc. B Biol. Sci* 282 (2015).
72. Rajendran P, dasan RB, esh KS, mar K. GC-MS Analysis of Phyto-Components in Raw and Treated Sugarcane Juice. *Int. J. Curr. Microbiol. Appl. Sci.* 6 (2017) 51–61.
73. RANI J, KAPOOR M. Gas Chromatography-Mass Spectrometric Analysis and Identification of Bioactive Constituents of *Catharanthus Roseus* and Its Antioxidant Activity. *Asian J. Pharm. Clin. Res* 12 (2019): 461–465.
74. Rees JA, M, M, P, B. Novel antibacterial peptides isolated from a European bumblebee, *Bombus pascuorum* (Hymenoptera, Apoidea). *Insect Biochem Mol Biol* 27 (1997): 413–422.
75. Seiichi Hara, Yamakawa M. A novel antibacterial peptide family isolated from the silkworm, *Bombyx mori*. *Biochem J* 310 (1995): 651–656.
76. Sewify GH, Hamada HM, Alhadrami HA. In Vitro Evaluation of Antimicrobial Activity of Alimentary Canal Extracts from the Red Palm Weevil, *Rhynchophorus ferrugineus* Olivier Larvae. *Biomed Res Int* (2017).

77. Sheeja L, Lakshmi D, Bharadwaj S, Sajidha Parveen K. Anticancer activity of phytol purified from *Gracilaria edulis* against human breast cancer cell line (MCF-7). *Int J Curr Sci* 19 (2016): 36–46.
78. Sheela D, Uthayakumari F. GC-MS ANALYSIS OF BIOACTIVE CONSTITUENTS FROM COASTAL SAND DUNE TAXON – *SESUVIUM PORTULACASTRUM* (L.). *Biosci. Discov* 4 (2013): 47–53.
79. Shoge M, Garba S, Labaran S. Antimicrobial Activities of 1,2-benzenedicarboxylic acid, butyldecyl ester isolated from the seeds and pods of *Acacia nilotica* Linn. *Basic Res. J. Microbiol* 3 (2016): 08–11.
80. Slade EM, Riutta T, Roslin T, Tuomisto HL. The role of dung beetles in reducing greenhouse gas emissions from cattle farming. *Sci. Rep.* 6 (2016).
81. Sosa AA, Bagi SH, Hameed IH. Analysis of bioactive chemical compounds of *Euphorbia lathyris* using gas chromatography-mass spectrometry and fourier-transform infrared spectroscopy. *J. Pharmacogn. Phyther* 8 (2016): 109–126.
82. Srivastava R, Mukerjee A, Verma A. GC-MS analysis of phytochemicals in, PET ether fraction of *wrightia tinctoria* seed. *Pharmacogn J* 7 (2015): 249–253.
83. Sudhendra Karthi BS and AJAH. *Journal of Chemical , Biological and Physical Sciences* Efficacy of Methanolic Extract of a Marine Ascidian , *Lissoclinum bistratum* for Antimicrobial Activity (2016).
84. Swamy MK, Arumugam G, Kaur R, Ghasemzadeh A, Yusoff MM, Sinniah UR. GC-MS Based Metabolite Profiling, Antioxidant and Antimicrobial Properties of Different Solvent Extracts of Malaysian *Plectranthus amboinicus* Leaves. *Evidence-based Complement. Altern Med* (2017).
85. Tarasov S, Génier F. Innovative bayesian and parsimony phylogeny of dung beetles (coleoptera, scarabaeidae, scarabaeinae) enhanced by ontology-based partitioning of morphological characters. *PLoS One* 10 (2015).
86. Taylor R, Romaine I, Liu C, Murthi P, Jones P, Waterson A, Sulikowski G, Zwiebel L. Structure-Activity Relationship of a Broad-Spectrum Insect Odorant Receptor Agonist. *ACS Chem. Biol* 7 (2012): 1647–1652.
87. Thevissen K, Warnecke DC, François IEJA, Leipelt M, Heinz E, Ott C, Zähringer U, Thomma BPHJ, Ferket KKA, Cammue BPA. Defensins from Insects and Plants Interact with Fungal Glucosylceramides. *J. Biol. Chem* 279 (2004): 3900–3905.
88. Turillazzi S, Perito B, Pazzagli L, Pantera B, Gorfer S, Tancredi M. Antibacterial activity of larval saliva of the European paper wasp *Polistes dominulus* (Hymenoptera, Vespidae). *Insectes Soc* 51 (2004): 339–341.
89. Venkata raman B, M PS, LA S, B NR, A NVK, M S, TM R. *Academic Sciences Asian Journal of Pharmaceutical and Clinical Research. Asian J. Pharm. Clin. Res* 5 (2012): 99–106.
90. Vetterli SU, Zerbe K, Müller M, Urfer M, Mondal M, Wang SY, Moehle K, Zerbe O, Vitale A, Pessi G, Eberl L, Wollscheid B, Robinson JA. Thanatin targets the intermembrane protein complex required for lipopolysaccharide transport in *Escherichia coli*. *Sci Adv* 4 (2018).
91. Vilas AK, Amit HM. Bioassay-guided fractional isolation and identification of anti-mrsa compounds by gms from *tridax procumbens*. *World J. Pharm Pharm Sci* 4 (2015): 1022–1031.
92. Vizioli J, Richman A, Uttenweiler-Joseph S, Blass C, Bulet P. The defensin peptide of the malaria vector mosquito *Anopheles gambiae*: antimicrobial activities and expression in adult mosquitoes. *Insect Biochem*

Mol Biol 31 (2001): 241-248.

93. Wu Q, Patočka J, Kuča K. Insect antimicrobial peptides, a mini review. *Toxins (Basel)* 10 (2018): 1–17.

94. Yi HY, Chowdhury M, Huang YD, Yu XQ. Insect antimicrobial peptides and their applications. *Appl. Microbiol. Biotechnol* 98 (2014): 5807–5822.

95. Zha C, Wang C, Li A. Toxicities of selected essential oils, silicone oils, and paraffin oil against the common bed bug (hemiptera: Cimicidae) *J Econ Entomol* 111 (2018): 170–177.

96. Zhang Z-M, Wu W-W, Li G-K. Study of the Alarming Volatile Characteristics of *Tessaratoma papillosa* Using SPME-GC-MS. *J. Chromatogr Sci* 47 (2009): 291–295.



This article is an open access article distributed under the terms and conditions of the

[Creative Commons Attribution \(CC-BY\) license 4.0](https://creativecommons.org/licenses/by/4.0/)