


Research Article

Skin Transparency Improvement Effects of Rice Vitamin E Crude Extract

Yuhō Yamauchi^{†1}, Jingxiu Zhou^{†1}, Shiori Nakagawa¹, Yoshikazu Uchida², Toshio Nakamura^{*1}, Takuo Tsuno¹

Abstract

Background: Rice bran contains various active ingredients that have attracted attention as materials for use in health foods and cosmetics. Vitamin E has beneficial antioxidant effects and are used in cosmetics. We have developed a method for producing Rice Vitamin E crude extract (RVE) which contains bioactive ingredients, such as tocopherols, tocotrienols, and plant sterols from the by-product of rice bran oil extraction. In this study, we verified the effectiveness of rice bran ingredients by using cell culture assays and face application tests in humans.

Methods: The expression of cyclooxygenase-2 (COX-2) and endothelin-1 (EDN-1), both of which promote melanogenesis, was analyzed in HaCaT cells with or without RVE treatment. In the face application test, RVE-containing cream and placebo cream were each applied to half of the face twice a day, and melanin index measurement and VISIA image analysis were performed.

Results: In cell assays, RVE reduced COX-2 and EDN-1 expression levels. In the application test, the melanin index on the RVE treated side significantly decreased over time from the start of the test, and after 12 weeks, it decreased significantly compared to that with the placebo. In addition, the analysis of the visual analog scale (VAS) questionnaire, which evaluates the participant's perception of the skin condition, indicated improvements in texture and transparency brought on by RVE.

Conclusions: This study indicates that RVE increases skin transparency by reducing the amount of melanin through suppression of COX-2 and EDN-1 production and by improving texture and elasticity.

Keywords: Rice bran; *Oryza sativa*; Antioxidant; Tocopherols; Tocotrienols; Transparency

Introduction

Rice is cultivated in many countries worldwide, mainly in East and Southeast Asia. Rice bran, a non-edible component of rice, accounts for approximately 10% of its weight and is known to contain various functional ingredients [1]. Among the functional ingredients of rice bran are tocopherols and tocotrienols, which belong to the vitamin E group. Tocotrienols have approximately 50 times the antioxidant power of tocopherols but are found in small quantities in only certain foods. We have developed Rice Vitamin E crude extract (RVE), containing more than 3% each of tocopherols and tocotrienols, by removing fatty acids and concentrating unsaponifiable substances from the deodorized fraction, which is a by-product of the rice oil refining process. It is reported that tocotrienols or tocotrienol-rich fraction inhibit melanogenesis by modulating the expression of genes involved in

Affiliation:

[†]Tsuno Rice Fine Chemicals Co., Ltd, Ito-gun, Katsuragi-cho, Chonomachi 2283, Wakayama, Japan

²Department of Food Science & Nutrition, Convergence Program of Material Science for Medicine and Pharmaceuticals, and The Korean Institute of Nutrition Hallym University, Chuncheon, Republic of Korea.

*Corresponding author:

Toshio Nakamura, Tsuno Rice Fine Chemicals Co., Ltd, Ito-gun, Katsuragi-cho, Chonomachi 2283, Wakayama, Japan.

Citation: Yuhō Yamauchi, Jingxiu Zhou, Shiori Nakagawa, Yoshikazu Uchida, Toshio Nakamura, Takuo Tsuno. Skin Transparency Improvement Effects of Rice Vitamin E Crude Extract. *Journal of Food Science and Nutrition Research*. 6 (2023): 31-39.

Received: December 23, 2022

Accepted: April 03, 2023

Published: April 20, 2023

Statistical analyses

In this study, the values are shown as the mean \pm standard error, and only the age of the participants is shown as the mean \pm standard deviation. The measured values of the clinical test were set to 0 before application of the cream (0 week), and the degree of change after a certain period was determined. Comparisons between groups in the cell test and comparisons before and after application of the test cream in the clinical trial were analyzed using Dunnett's test; a risk rate of 5% or less was considered a significant difference. In addition, the clinical test results were compared between the placebo-treated side and RVE-treated side and analyzed using Student's t-test. Cases with a risk rate \leq 5% were found to have a significant difference. MEPHAS, a statistical analysis program for pharmaceutical data provided by the Center for Genetic Information Experiments at Osaka University, was used in this study.

Results

Expression levels of COX-2 and EDN-1 in cultured epidermal keratinocyte cells HaCaT

RT-qPCR was used to validate the effects of RVE on COX-2 and EDN-1 expression. COX-2 expression was significantly decreased in cells treated with 0.001% and 0.01% RVE compared with that in untreated cells (Figure 2a). A significant decrease in EDN-1 expression was also observed with the 0.001% and 0.01% RVE treatments (Figure 2b).

The expression levels of COX-2 (a) and EDN-1(b) in HaCaT examined by real-time PCR. The expression level of each gene is given as a ratio to that of GAPDH, and the expression of RVE-untreated samples is denoted as 1. Data are presented as the mean \pm SE. (n = 4) Data were analyzed by Dunnett's test. *; $p < 0.05$ (vs. RVE-untreated samples)




Figure 2: Related factors of melanin synthesis

Results of participant-grouping

The study design for topical application of RVE is presented in table 3. Twenty-five participants were assessed for eligibility, and three did not meet the inclusion criteria. Twenty-two participants were divided into two groups: one group applied RVE cream on the right side of the face, and the other group applied RVE cream on the left side. There was no significant difference between the initial values of the RVE-treated- and placebo-treated sides for any of the measurement parameters. The mean age of the participants at the start of the study was 37.5 ± 11.7 years. One participant dropped out during the test period, and the final analysis was conducted on 21 participants.

Melanin index

The melanin index measured by Mexameter MX18 showed a significant decrease from the initial value only at 8th week in the placebo group ($p < 0.05$) and in the RVE group at 4th, 8th, and 12th week ($p < 0.05$). The decrease from the initial measurement at 12th week was greater in the RVE group than that in the placebo group ($p < 0.05$) (Figure 3).

Changes from the 0th week in melanin index. The dotted line shows placebo treated sides and the solid line shows the RVE treatment sides. Data are presented as the mean \pm SE. Data were analyzed by Dunnett's test. *; $p < 0.05$ (vs. 0th week RVE), #; $p < 0.05$ (vs. 0th week placebo), †; $p < 0.05$ (vs. placebo treated sides).

Results of VISIA analysis

Scores of the spots and pores were analyzed using VISIA evolution. There were no significant differences in the measurements between the initial and later stages and between the placebo and RVE sides (Figure 4).




Figure 6: Result of VAS questionnaire

side showed a significant score increase from the initial value at 8th week ($p < 0.05$) (Figure 6c).

Changes from 0th week in the score of texture (a), elasticity (b), and transparency (c) by VAS questionnaire. The dotted line shows placebo treated sides and the solid line shows the RVE treatment sides. Data are presented as the mean \pm SE. * ; $p < 0.05$ (vs. 0th week RVE), † ; $p < 0.05$ (vs. placebo treated sides).

Discussion

We examined the cosmetic effect of RVE on human skin by analyzing the gene expression of melanin production-promoting factors in cultured cells and the amount of melanin, spots, and pore scores in a half-face application test. In cultured cells, reduced expression levels of COX-2 and EDN-1, promoters of tyrosinase activity, was observed. We also found that RVE reduced the amount of melanin on human skin. There was no significant difference in the VISIA blemish score, but the decrease in the RVE treated side was greater than that in the placebo side at 4 and 8 weeks. RVE is thought to reduce melanin and spots because the vitamin E group in RVE contains tocotrienol, which is reported to have a strong antioxidant effect and removes reactive oxygen species generated by UV rays, thereby reducing COX-2 and EDN-1 synthesis.

The pore score, analyzed by VISIA, did not decrease due to RVE application. In this study, we included the pore score as a secondary outcome in anticipation of a reduction in pore darkening owing to the inhibition of lipid peroxide production by the antioxidant action of RVE. A once-daily application of a comprehensive anti-oxidant product containing vitamins C and E, among others, was reported to improve pores in men's skin as assessed by visual inspection by an expert [12]. On the other hand, in a study investigating the influence of astaxanthin-containing beverages, which have an anti-lipid peroxidation effect on the skin, positive effects were seen in skin texture and moisture content, but no effect was seen in pores analyzed by VISIA [13]. In the VISIA analysis, the score is calculated from the number and area of pores, and therefore, it is not possible to strictly evaluate changes in darkening due to oxidation of sebum in pores. Since the darkening of pores is also related to a decrease in skin transparency, the VAS questionnaire showed an improvement in transparency; therefore, we presumed that there was no effect on the conspicuity of pores.

Regarding skin elasticity, the R5 value, which is the net elasticity, and the R7 value, which is the recovery after deformation, tended to increase from the initial measurement values on the RVE side compared to those on the placebo side, although they were not statistically significant. The tendency

to increase observed in R5 and R7 is consistent with the VAS questionnaire results, which showed a significantly greater change from the initial value in the 4th week in the RVE treated side than that in the placebo side. Similar to RVE, the phytosterol ester extract, which is rich in plant sterol esters, showed the effect of promoting type 3 collagen production in fibroblasts in our laboratory. In addition, since vitamin E has an antioxidant effect, it is thought to repair the structure of the dermis damaged by oxidative stress. These findings suggest that the plant sterol esters and vitamin E contained in RVE contribute to the improvement of elasticity.

In addition to the elasticity mentioned above, the VAS questionnaire showed that the application of RVE improved the texture and transparency. It has been reported that the fineness of texture and the amount of melanin are involved in the transparency of the skin [14]. Another study that systematically analyzed the interaction between clarity and skin physiology found that “red spot count” and “skin smoothness” and “yellowness or lightness in skin color” as key characteristics that dictate skin transparency [15]. In addition, it has been reported that the carbonylation of the stratum corneum protein is involved in the yellowing of the skin and the loss of transparency [16,17]. Carbonylated protein has been used as a measure of protein oxidation [18,19], and it should be considered that this might be improved by RVE containing vitamin E group with antioxidative effects. The transparency of the skin is considered to be a major factor in beautiful skin, so there should be a high demand for its improving effect.

In this study, RVE was shown to have melanin-reducing, elasticity-increasing, and texture-improving actions and has been found to be a valid cosmetic raw material with multiple effects of improving skin transparency.

Acknowledgements

We would like to thank Editage (www.editage.com) for English language editing.

References

- Ghosh M. Review on recent trends in rice bran oil processing. *J Am Oil Chem Soc* 84 (2007): 315-324.
- Michihara A, Ogawa S, Kamizaki Y, et al. Effect of delta-tocotrienol on melanin content and enzymes for melanin synthesis in mouse melanoma cells. *Biol Pharm Bull* 33 (2010): 1471-1476.
- Makpol S, Jam FA, Rahim NA, et al. Comparable down-regulation of TYR, TYRP1 and TYRP2 genes and inhibition of melanogenesis by tyrostat, tocotrienol-rich fraction and tocopherol in human skin melanocytes improves skin pigmentation. *Clin Ter* 165 (2014): e39-e45.
- Eo S H, Kim S J. Resveratrol-mediated inhibition of cyclooxygenase-2 in melanocytes suppresses melanogenesis through extracellular signal-regulated kinase 1/2 and phosphoinositide 3-kinase/Akt signalling. *European Journal of Pharmacology* 860 (2019): 172586.
- Hachiya A, Kobayashi A, Ohuchi A, et al. The paracrine role of stem cell factor/c-kit signaling in the activation of human melanocytes in ultraviolet-B-induced pigmentation. *J. Investig. Dermatol* 116 (2001): 578-586.
- Delinasios GJ, Karbaschi M, Cooke MS, et al. Vitamin E inhibits the UVAI induction of “light” and “dark” cyclobutane pyrimidine dimers, and oxidatively generated DNA damage, in keratinocytes. *Sci Rep* 8 (2018): 423.
- Wu CM, Cheng YL, Dai YH, et al. alpha-Tocopherol protects keratinocytes against ultraviolet A irradiation by suppressing glutathione depletion, lipid peroxidation and reactive oxygen species generation. *Biomed. Rep* 2 (2014):419-423.
- Gao LX, Bian C, Wu Y, et al. Label-free electrochemical sensor to investigate the effect of tocopherol on generation of superoxide ions following UV irradiation. *J Biol Eng* 12 (2008): 17.
- Loizou S, Paraschos S, Mitakou S, et al. Chios mastic gum extract and isolated phytosterol tirucallol exhibit anti-inflammatory activity in human aortic endothelial cells. *Exp Biol Med* 234 (2009): 553–61.
- Adachi H, Murakami Y, Hotta N, et al. Investigation of the skin aging mechanism focusing on stratifin. *J Soc Cosmet Chem Jpn* 49 (2015): 211-217.
- Nakamura-Utsunomiya A, Tsumura M, Okada S, et al. Downregulation of endothelial nitric oxide synthase (eNOS) and endothelin-1 (ET-1) in a co-culture system with human stimulated X-linked CGD neutrophils. *PLoS One* 15 (2020): e0230665
- Sikora BC, Wortzman M, Nelson DB, et al. A pilot study evaluating the efficacy and tolerability of a comprehensive, hydrating topical antioxidant developed specifically for men. *J Cosmet Dermatol* 20 (2021): 2816-2823.
- Tsukahara H, Matsuyama A, Abe T, et al. Effects of intake of astaxanthin contained drink on skin condition. *Jpn J Complementary Altern Med* 13 (2016): 57-62.
- Masuda Y, Kunizawa N, Takahashi M. Methodology for evaluation of skin transparency and the efficacy of an essence that can improve skin transparency. *J Soc Cosmet Chem* 39 (2005): 201-208.
- Nakamura R, Itai E, Uehara S, et al. Development of Mathematical Model of Skin Translucency. *J. Soc Cosmet. Chem. Jpn* 54 (2020): 4.
- Masaki, H, Mizutani T, Ogawa N, et al. Carbonylated

- proteins contribute to the darkness around facial pores. *J. Dermatol. Sci* 89 (2008): 299-307.
17. Ichiro Iwai, Tomohiro Kuwahara, Tetsuji Hirao. Decrease in the Skin Transparency Induced by Protein Carbonylation in the Stratum Corneum. *J. Soc. Cosmet. Chem. Jpn* 43 (2008): 1.
18. CS Sander, H Chang, S Salzmann, et al. Thiele Photoaging is associated with protein oxidation in human skin in vivo *J. Invest. Dermatol* 118 (2022): 618-625.
19. Laurent C, LeDuc A, Pottier I, et al. Dramatic Increase in Oxidative Stress in Carbon-Irradiated Normal Human Skin Fibroblasts. *PLoS ONE* 8 (2013): e85158.