



Research Article

The Spermine Phosphate-Bound Cyclooctaoxygen Sodium Epigenetic Shell of Euchromatin DNA Is Destroyed by the Epigenetic Poison Glyphosate

Andreas J Kesel*

Abstract

Oxygen exists in two gaseous (dioxygen and ozone) and six solid allotropic modifications. An additional allotropic modification of oxygen, the cyclooctaoxygen, was predicted to exist in 1990. The first synthesis and characterization of cyclooctaoxygen as its sodium crown complex, isolated in the form of three cytosine nucleoside hydrochloride complexes, was reported in 2016. The sperminium hydrogen phosphate/cyclooctaoxygen sodium complex was calculated to cover the actively transcribed regions (2.6%) of bovine lymphocyte interphase genome. Cyclooctaoxygen seems to be naturally absent in hypoxia-induced highly condensed chromatin, taken as a model for eukaryotic metaphase/anaphase/early telophase mitotic chromatin. Hence, it is proposed that the cyclooctaoxygen sodium-bridged sperminium hydrogen phosphate and selenite coverage serves as an epigenetic shell of actively transcribed gene regions in eukaryotic ‘open’ euchromatin DNA. Cyclooctaoxygen sodium-bridged sperminium hydrogen selenite was calculated to serve as a marker shell component at ATG start codons in human euchromatin DNA mRNA genes, both at the translation initiation triplet and at 5'-untranslated region upstream ATGs. The total herbicide glyphosate (ROUNDUP®) and its metabolite (aminomethyl)phosphonic acid (AMPA) are proved to represent ‘epigenetic poisons’, since they both selectively destroy the cyclooctaoxygen sodium complex. This definition is of reason, since the destruction of cyclooctaoxygen is certainly sufficient to bring the protection shield of human euchromatin into collateral epigenetic collapse. The total herbicide glyphosate and its environmental metabolite (aminomethyl)phosphonic acid (AMPA) can be associated *in vitro* with catalytic deterioration of eukaryotic euchromatin genetic information. The epigenetic shell of eukaryotic euchromatin is susceptible to decay induced by catalytic epigenetic poisons threatening eukaryotic genomic heritage.

Keywords: AMPA; Cyclooctaoxygen; DNA; Glyphosate; Selenium; Spermine

Introduction

In 1677 *Antoni van Leeuwenhoek* discovered [1] the characteristic crystals of spermine phosphate (spermine \times 2 H₃PO₄ \times 6 H₂O) [2] in matured native human semen (Figure 1). Since then, there was collected conclusive evidence that human chromosomal DNA is closely associated with spermine phosphate [2]. The interaction of oxygen species with DNA has until recently only been focused on oxidative DNA damage and its pathophysiological consequences [3]. In 2015 *Kirmes et al.* reported an unprecedented interaction of eukaryotic chromatin DNA structure with atmospheric oxygen partial pressure [4]. Under switching to hypoxic conditions (1% O₂, 5% CO₂, 94% N₂) the murine cardiomyocyte

Affiliation:

Chammünsterstr. 47, D-81827 München, Bayern/
Bavaria, Germany

*Corresponding author:

Andreas J Kesel, Chammünsterstr. 47, D-81827
München, Bayern/Bavaria, Germany.

Citation: Andreas J Kesel. The Spermine Phosphate-Bound Cyclooctaoxygen Sodium Epigenetic Shell of Euchromatin DNA Is Destroyed by the Epigenetic Poison Glyphosate. *Archives of Clinical and Biomedical Research* 7 (2023): 171-190.

Received: February 05, 2023

Accepted: February 15, 2023

Published: March 16, 2023

HL-1 cell chromatin rendered itself highly condensed, accompanied by redistribution of the polyamine pool (mainly spermine and spermidine) to the nucleus [4]. In 2016 Kesel *et al.* showed [5] that eukaryotic single-stranded DNA (ssDNA) binds a new allotropic form of oxygen, the cyclooctaoxygen (cyclo-O₈), in form of its sodium (Na⁺) complex (cyclo-O₈-Na⁺), especially when in coordination to spermine phosphate (sperminium hydrogen phosphate) [5]. A model for a logically resulting first epigenetic shell of eukaryotic *in vivo* DNA was proposed [5]. Also a partial substitution of the sperminium (C₁₀H₃₀N₄⁴⁺)-bound monohydrogen phosphate (HPO₄²⁻) anions by hydrogen selenite (HSeO₃⁻) anions was postulated during these investigations [5], thereby providing an explanation for the well-known, but ‘mysterious’ [6], augmenting effects of the polyamine spermine [6] and the essential micronutrient selenium [5] on eukaryotic genome integrity and chromosomal DNA stability.

Materials and Methods

Materials

The following materials were utilized: **NC** (C₉H₁₄ClN₃O₅ × 2 ClNaO₈ × 2 H₂O × ½ C₃H₆O) = [tetrakis(β-D-cytidin-N³-ium)(octoxocane-κ⁴O¹,O³,O⁵,O⁷)sodium(5+) pentachloride dihydrate hemiacetonate (*M* = 1,370.22 g/mol) [5]; **RC** (C₉H₁₄ClN₃O₅ × 2 HClNa₂O₁₇ × ¼ C₃H₆O) = β-D-cytidine hydrochloride – μ-chloro(μ-hydroxy)bis(octoxocane-κ⁴O¹,O³,O⁵,O⁷)disodium (1:2) × 0.25 acetone (*M* = 1,003.06 g/mol) [5]; potassium iodide (KI) *puriss. p.a.*, reagent, ISO, reagent, Ph.Eur., *w* (*m/m*) ≥ 99.5% (*M* = 166.00 g/mol) {Sigma-Aldrich Corp., St. Louis, MO, USA; pH 6.0–9.0 [*θ* = 20 °C, 5% (*m/m*) in H₂O], loss on drying ≤ 0.2% (*θ* = 105 °C), total nitrogen (N) ≤ 0.001%, heavy metals (as Pb) ≤ 0.0005%, iodate (IO₃⁻) ≤ 2 mg/kg}; starch *puriss. p.a.*, from potato, reagent, ISO, reagent, Ph.Eur., soluble [Sigma-Aldrich Corp., St. Louis, MO, USA; pH 6.0–7.5, loss on drying ≤ 13% (*θ* = 105 °C), sulfated ash ≤ 0.5%, substances reducing *Fehling* solution (as maltose) ≤ 0.7 %]; (aminomethyl)phosphonic acid (AMPA) (CH₆NO₃P, *M* = 111.04 g/mol) [Sigma-Aldrich Corp., St. Louis, MO, USA, Lot: MKBX8824V; *w* (*m/m*) = 98.5% (titration), carbon (C) 10.9%, nitrogen (N) 12.6%]; *N*-(phosphonomethyl)glycine (glyphosate) (C₃H₈NO₅P, *M* = 169.07 g/mol) [Sigma-Aldrich Corp., St. Louis, MO, USA, Lot: MKBX1937V; *w* (*n/n*) > 99% (TLC), carbon (C) 21.3%, nitrogen (N) 8.2%, mp 230 °C (dec.)]; deuterated chloroform (CDCl₃) [eurisotop®, Saint Aubin cedex, France, Lot: D007H, W2631; 99.80% D, H₂O < 0.01%, stored over molecular sieve 3Å]; molecular sieve 3Å (0.3 nm, zeolithe, metal-aluminosilicate) [AppliChem GmbH, Darmstadt, Germany; Lot: 5H002478; water absorbency ≥ 20% (24 h, 80% humidity)]; elemental iodine I₂ (*M* = 253.81 g/mol) (iodum resublimatum Ph.Eur. 7.0) [Caesar & Loretz (Caelo) GmbH, Hilden, Germany, Lot: 122890; *w* (*m/m*) = 99.9% (titration), non-volatile matter ≤ 0.1%, Br⁻ and Cl⁻ ≤ 250 ppm]. L-Ascorbic acid (vitamin C)

Ph.Eur. 7.0 (*M* = 176.12 g/mol) was purchased from a local pharmacy store (Friedens-Apotheke, München-Trudering, Germany). Glyphosate monosodium salt (glyphosate-Na, C₃H₇NNaO₅P, *M* = 191.05 g/mol) was purchased in form of ROUNDUP® GRAN 15,5 g granules [Monsanto Europe S.A., Antwerp, Belgium, Lot: T346 (prod. date 2011/12/12); contains 475 g/kg glyphosate-Na, *w*_{glyphosate-Na} (*m/m*) = 47.5%, inert carrier material 33.5%, water and excipients 19%].

Calculation of the Genomic Coverage by the Cyclooctaoxygen Sodium-Bridged Spermine Phosphate Epigenetic Shell of Interphase DNA in Bovine Lymphocytes

The average volume of a human blood lymphocyte was taken as 206 f(emto)l(iter) [10⁻¹⁵ liter = (μm)³] [7, 8] after technical correction [8] of the published value (210 fl) [7]. The published average volume of a bovine blood lymphocyte (256 fl) [9] was taken as 214 fl after introducing a technical correction factor of *f* = 0.834 [8]. The reference genome size of *Bos taurus* (Hereford breed) was taken as 2,670,139,648 bp (RefSeq assembly accession number GCF_000003055.6) [10]. This genome showed 41.89% GC content [10]. The theoretical intracellular concentration of the sperminium hydrogen phosphate/cyclo-O₈-Na⁺ complex required to cover all triplets of the dsDNA genome in a blood lymphocyte of *B. taurus* was calculated: [214 fl × 6.022140857 × 10²³ mol⁻¹]⁻¹ × 2,670,139,648 × 2 × (3)⁻¹ = 13.8127 mM. The coverage of *B. taurus* genome by the sperminium hydrogen phosphate/cyclo-O₈-Na⁺ complex was calculated (mean ± s.d.: 2.6208 ± 0.4953%) from the published [11] fractions of spermine bound to dsDNA: 421 μM × [13.8127 mM]⁻¹ = 3.0479% (in presence of 2 mM Mg²⁺ and 100 mM K⁺), 287 μM × [13.8127 mM]⁻¹ = 2.0778% (2 mM Mg²⁺, 150 mM K⁺), and 378 μM × [13.8127 mM]⁻¹ = 2.7366% (10 mM Mg²⁺, 100 mM K⁺). The number of base pairs for the number of protein-coding exons in *B. taurus* genome (49,107) [10] was calculated with the median of amino acid residues/gene (468) [10]: 49,107 × 3 bp × 468 = 68,946,228 bp (2.5821% of *B. taurus* genome).

Calculation of the Spermine Coverage of Highly Condensed Mitotic Metaphase DNA in HeLa S3 Cells

The average effective molecular mass of dGp/dCp was calculated as *M* = 309.19 g/mol, of dAp/dTp as *M* = 308.70 g/mol. The reference *Homo sapiens* genome size was taken as 3,099,441,038 bp (RefSeq assembly accession number GCF_000001405.40) [12]. This genome showed 40.5% GC content [12]. The molecular mass of this human genome dsDNA was calculated: [(0.405 × 309.19 g/mol) + (0.595 × 308.70 g/mol)] × [3,099,441,038 bases (b) × 2] = 1.191482507 × 10¹² g/mol. The HeLa cell genome [13] was anticipated as 76 chromosomes (hypertriploid) + 22 abnormal chromosomes [13,14]. The chromosomal DNA

size of HeLa metaphase chromatin dsDNA was calculated as 19,539,129,390 bp with $M = 1.207127414 \times 10^{13}$ g/mol from published karyotyping [14]. The content of spermine in HeLa S3 cell metaphase chromatin was taken as 135.9 ± 16.1 pmol/ μ g DNA [15]. This was transformed into 135.9 ± 16.1 p(ico)mol (10^{-12} mol) spermine/82.84129651 z(epto)mol (10^{-21} mol) dsDNA = 1,640,486,155 (molecules spermine/dsDNA). Since one spermine molecule is assumed to cover six bp (in the pure spermine form of A-DNA duplex [16] and Z-DNA duplex [17]), this corresponds to a coverage of $1,640,486,155 \times 6 \times [19,539,129,390]^{-1} = 50.3754\%$ of HeLa S3 cell metaphase chromatin dsDNA genome by spermine. The content of spermidine in HeLa S3 cell metaphase chromatin was taken as 116.1 ± 11.8 pmol/ μ g DNA [15]. This was transformed into 116.1 ± 11.8 p(ico)mol (10^{-12} mol) spermidine/82.84129651 z(epto)mol (10^{-21} mol) dsDNA = 1,401,474,927 (molecules spermidine/dsDNA). Since one spermidine molecule is assumed to cover six bp (in the pure spermidine form of Z-DNA duplex [18]), this corresponds to a coverage of $1,401,474,927 \times 6 \times [19,539,129,390]^{-1} = 43.0359\%$ of HeLa S3 cell metaphase chromatin dsDNA genome by spermidine. Taken together, the polyamine (spermine/spermidine ratio 1.17) coverage of HeLa S3 cell metaphase chromatin dsDNA is $50.3754\% + 43.0359\% = 93.4113\%$.

Calculation of the Polyamine Coverage of Maximally Condensed Mitotic Late Anaphase/Early Telophase DNA in Murine Cryptal Enterocytes

The phosphorus (P) content of female *Mus musculus* strain C3H/HeJ cryptal enterocytic mitotic (late anaphase/early telophase) chromatin was taken as 298.5 ± 17.3 mmol (P)/kg [19]. The average effective molecular mass of one nucleotide unit (dGp/dCp and dAp/dTp) in *M. musculus* C3H/HeJ genome (2,701,131,316 bp) [20] was calculated with the GC content of 42.82% [20]: $M = 308.91$ g/mol. With the formula $w(P) = (n_p) \times M^{-1}$ (n_p , number of P atoms *pro* formula unit; M , molecular weight of formula unit) the phosphorus content $w(P)$ of murine mitotic chromatin was calculated: 3,237.19 mmol (P)/kg (free dsDNA). The value for complete complexation with one spermine tetracation/six nucleotides is 2,912.86 mmol (P)/kg (spermine tetracation-complexed ssDNA), and for complete complexation with one spermidine trication/six nucleotides is 2,997.41 mmol (P)/kg (spermidine trication-complexed ssDNA). Taking the arithmetic mean of the published [11, 21] spermine/spermidine ratio 0.85, bound to rat liver DNA and RNA (2 mM Mg^{2+} , 150 mM K^+) [11], or in *Sprague-Dawley* rat liver nuclei [21], the theoretical value for complete complexation is $\{(2,912.86 \times 0.85) + [(2,997.41 \times (2 - 0.85))]\} \times (2)^{-1} = 2,961.48$ (P)/kg (spermine tetracation/spermidine trication-complexed ssDNA). Phosphatidylcholine in its dipalmitoyl ($C_{16:0}$) lipid composition ($M = 734.04$ g/mol) is the main constituent (60.8 \pm 1.3%) of the phospholipid fraction in rat liver chromatin

[22]. This molecular weight was corrected to an average effective molecular mass $M = 765.03$ g/mol, based on the fractions [22] and lipid compositions [22] of phospholipids (phosphatidylcholine, phosphatidylethanolamine, phosphatidylserine, phosphatidylinositol, sphingomyelin) in *Sprague-Dawley* rat liver chromatin, to yield the reference value for the (P) content of phospholipid 1,307.13 mmol (P)/kg. The fractions of DNA ($32.0 \pm 4.1\%$), RNA ($5.1 \pm 1.6\%$), protein ($62.6 \pm 3.8\%$), and phospholipid ($0.2 \pm 0.1\%$) in *Sprague-Dawley* rat liver chromatin were taken as published [22], and applied on the murine mitotic chromatin nucleic acids (DNA + RNA). The published [19] value was corrected for the chromatin-bound cation (Na^+ , K^+ , Mg^{2+} , Ca^{2+}) [19], protein and phospholipid [22] content: $\{[841.0 \text{ mmol } (Na^+, K^+, Mg^{2+}, Ca^{2+})/kg + 298.5 \text{ mmol } (P)/kg] \times 100 \times (37.1)^{-1}\} + [0.2 \times (100)^{-1} \times 1,307.13 \text{ mmol } (P)/kg] = 3,074.04$ mmol (P)/kg (ssDNA + ssRNA) in mitotic late anaphase/early telophase chromatin. The six nucleotide coverage (ssDNA + ssRNA) by the polyammonium cations was calculated: $(3,237.19 - 3,074.04) \times (3,237.19 - 2,961.48)^{-1} \times 100\% = 59.1745\%$. The genomic dsDNA six bp coverage by the polyammonium cations, corrected for the nuclear RNA content, was calculated: $59.1745\% \times [32.0 \times (37.1)^{-1}] \times 2 = 102.0800\%$. This corresponds to a coverage of 5.8777 bp murine cryptal enterocytic mitotic (late anaphase/early telophase) chromatin dsDNA by one polyammonium cation (spermine/spermidine ratio 0.85). The six nucleotide coverage of nuclear RNA by the polyammonium cations was calculated: $59.1745\% - [102.0800\% \times (2)^{-1}] = 8.1345\%$, corresponding to a relative six nucleotide coverage of nuclear RNA: $8.1345 \times (5.1)^{-1} \times 100\% = 159.5000\%$. This corresponds to a coverage of 3.7618 nucleotides of nuclear RNA by one polyammonium cation (spermine/spermidine ratio 0.85).

Calculation of the Hydrogen Selenite ($HSeO_3^-$) Coverage of Human Euchromatin DNA Specifically at ATG Start Codon Sequences

The reference *H. sapiens* genome size was taken as 3,099,441,038 bp (RefSeq assembly accession number GCF_000001405.40) [12]. The number of genes and pseudogenes in *H. sapiens* genome (59,265) was taken as published [12]. The median of the number of amino acid (aa) residues for a protein encoded by the actively transcribed part of the human genome (euchromatin) was calculated with the number of exon transcripts (394,894) [12] exhibiting the median length 143 b [12], and the number of single-exon (none intron to be spliced) transcripts (1,770) [12] exhibiting the median length 1,595 b [12]. The sum of the median exon transcript length and the median single-exon transcript length $[(394,894 \times 143 \text{ b}) + (1,770 \times 1,593 \text{ b})] = 59,289,452$ b (number of bases coding for euchromatin mRNA) was divided by the number of genes (41,989) [12] to yield the median length of euchromatin mRNA transcripts

$[(394,894 \times 143 \text{ b}) + (1,770 \times 1,593 \text{ b})] \times (41,989)^{-1} = 1,412 \text{ b}$. This value was divided by three (as to the triplet nature of the genetic code) to give the median of aa residues/gene of human euchromatin $[(394,894 \times 143 \text{ b}) + (1,770 \times 1,593 \text{ b})] \times (41,989)^{-1} \times (3)^{-1} = 470.6745 \text{ aa}$. The percentage of actively transcribed regions in the human genome was calculated: $470.6745 \text{ aa} \times 3 \times 59,265 \times (3,099,441,038 \text{ bp})^{-1} = 2.7000\%$ of *H. sapiens* genome. The theoretical intracellular concentration of the sperminium hydrogen phosphate/cyclo- $\text{O}_8\text{-Na}^+$ complex required to cover all triplets of the dsDNA genome in a blood lymphocyte of *H. sapiens* was calculated: $[206 \text{ fl} \times 6.022140857 \times 10^{23} \text{ mol}^{-1}]^{-1} \times 3,099,441,038 \text{ b} \times 2 \times (3)^{-1} = 16.6561 \text{ mM}$. The average volume of a human blood lymphocyte was taken as 206 fl [femtoliter = $(\mu\text{m})^3$] [7, 8] after technical correction [8] of the published value (210 fl) [7]. The optimal human blood serum selenium (as Se, $M = 78.96 \text{ g/mol}$) level of $110 \mu\text{g/l}$ (110 ng/ml , this serum level corresponds to a concentration of $1.3931 \mu\text{M}$) was taken from literature as concentration for optimal selenoprotein expression [23, 24]. The percentage of selenium (as selenite) concentration accumulated intracellularly in the cytosol from extracellular medium was taken from literature [25] [as 57.6% uptake into the cytosol of primary rat hepatocytes (1 min time, $10 \mu\text{M Na}_2^{75}\text{SeO}_3$ extracellularly)]. The median number of aa-coding triplets required for one triplet to be covered with the sperminium selenite/cyclo- $\text{O}_8\text{-Na}^+$ complex in interphase euchromatin (in form of hydrogen selenite HSeO_3^-) was calculated by: $(16.6561 \text{ mM} \times 2.7000\% \times 78.96 \text{ g/mol}) \times (110 \mu\text{g/ml} \times 0.576)^{-1} = 560.4400 \text{ aa}$. The number of 5'-untranslated region (5'-UTR) upstream open reading frames (uORFs) upstream start codons (uATGs transcribed to uAUGs) in the 5'-UTR of human mRNA genes was calculated from literature [26] as the number of transcribed human genes with at least one uATG (11,670) divided by the number of human mRNA genes with annotated 5'-UTR (23,775): $11,670 \times (23,775)^{-1} = 49.0852\%$. That means that 49% of human mRNA genes contain at least one uATG 5'-UTR start codon [26] [the downstream 3'-untranslated region (3'-UTR) contains none ATG start codon]. The total number of human uATGs/5'-UTR of human mRNA genes (16,504) was published (35,599) [27]. This gives the number of human uATGs/5'-UTR: $35,599 \times (16,504)^{-1} = 2.1570 \text{ uATGs/5'-UTR (human)}$. If this number is multiplied by the percentage of human mRNA genes containing at least one uATG 5'-UTR start codon [26], the number of hydrogen selenite HSeO_3^- molecules (within the sperminium hydrogen selenite/cyclo- $\text{O}_8\text{-Na}^+$ complex in interphase euchromatin) bound to human mRNA genes can be calculated: $2.1570 \times 49.0852\% = 1.0588 \text{ HSeO}_3^-/5\text{'-UTR of mRNA gene (human)}$. Accordingly, together with the transcribed and translated gene open reading frame initiation codon ATG encrypted in the Kozak consensus sequence 5'-ACCATGG-3' [28], the number of

human mRNA gene-incorporated hydrogen selenite HSeO_3^- molecules (within the sperminium hydrogen selenite/cyclo- $\text{O}_8\text{-Na}^+$ complex in interphase euchromatin) can be calculated: $1 (\text{HSeO}_3^-/\text{initiation ATG}) + 1.0588 (\text{HSeO}_3^-/5\text{'-UTR}) = 2.0588 \text{ HSeO}_3^-/\text{mRNA gene (human)}$. The median length of human mRNA transcripts ($3,560 \text{ b}$) [12] can be transformed into the median aa length (triplets) of human translated genes (as to the triplet nature of the genetic code): $3,560 \text{ b} \times (3)^{-1} = 1,186.6667 \text{ aa}$. If divided by the number of human mRNA gene-incorporated hydrogen selenite HSeO_3^- molecules, the number of triplets required for the covering by one hydrogen selenite HSeO_3^- molecule (within the sperminium selenite/cyclo- $\text{O}_8\text{-Na}^+$ complex in interphase euchromatin) can be calculated: $1,186.6667 \text{ aa} \times \{2.0588 [\text{HSeO}_3^-/\text{mRNA gene (human)}]\}^{-1} = 576.3978 \text{ aa}$. From this value the optimal human blood serum selenium (as Se, $M = 78.96 \text{ g/mol}$) level required for covering of all ATG codons in human mRNA genes (the downstream 3'-UTR contains none ATG start codon) was calculated: $(16.6561 \text{ mM} \times 2.7000\% \times 78.96 \text{ g/mol}) \times (576.3978 \text{ aa} \times 0.576)^{-1} = 106.9546 \mu\text{g/l (ng/ml; } 1.3545 \mu\text{M)}$.

Calculation of the Apparent Acid Dissociation Constant of the Human Genome DNA

The intrinsic pK_a of one isolated phosphodiester of DNA is 1.29, this is the pK_a ($\vartheta = 25 \text{ }^\circ\text{C}$) of dimethyl phosphate [29]. The reference *H. sapiens* genome size was taken as 3,099,441,038 bp (RefSeq assembly accession number GCF_000001405.40) [12]. The apparent (effective) $\text{pK}'_{a,\text{HG}} = 2.5601$ ($\vartheta = 25 \text{ }^\circ\text{C}$) of the haploid human genome was calculated (Figure 2A) according to the method of *Katchalsky & Gillis* [30], as based on the theoretical considerations of *Kuhn & Kuhn* [31]. The length rise of one B-DNA repeating unit (helix rise/bp) in solution (in the crystal: 0.336 nm [32]) was taken as: $b = 0.334 \times 10^{-7} \text{ cm}$ ($0.334 \pm 0.01 \text{ nm}$ [32]). The h_0 was calculated by: $0.334 \text{ nm} \times 3,099,441,038 = 1.03521331 \text{ m}$ (= theoretical length of *H. sapiens* dsDNA haploid genome as linear thread). The h_v was calculated by: $0.5 \times (0.334 \text{ nm} + 0.5798 \text{ nm}) \times 3,099,441,038 = 1.41613461 \text{ m}$ (theoretical length of half-neutralized *H. sapiens* dsDNA haploid genome as linear thread) (Figure 2A). The intermediate term ($n - 1$) was calculated as 0.4763 with $\lambda = 2.0321$ and $\lambda' = 0.0321$ (Figure 2A). The 0.5798 nm pitch in one relaxed (fully neutralized) dsDNA repeating unit was calculated as statistic arithmetic mean $d_0 = 579.80 \text{ pm}$ from the interphosphorus distance d ($5\text{'-P, } 5\text{'-P}$) = $739.25 \pm 14.37 \text{ pm}$ ($d \pm \text{s.d.}$; $n = 4$) in the fully ionized single-stranded hexanucleotide d(ApApApApApAp) (molecular modeling software: ACD/Chem Sketch version 12.01 with integrated ACD/3D Viewer, processed with Mercury 3.1 version 3.1.1), minus the typical phosphorus-oxygen distance d (P, O) = 159.45 pm [33], by the formula: $d_0 = 739.25 \text{ pm} - 159.45 \text{ pm} = 579.80 \text{ pm}$.

A *Observationes D. Anthonii Lewenhoeck, de Natis è semine genitali Animalculis.*

Nec non Auctoris harum Transactionum Responſa.

Observatoris Epistola Honor.iff. D. D. Vicecomiti Brauncker, Latinè conscripta; Dat. Nov. 1677. quam ipsissimis tunc transmissis verbis inferendam Auctor censuit.

Nobilissimè Vir,

Ultima ad Vestram Nobilitatem data littera praterit mensis decimo sexto, quamvis jam Nob. Vestra utilissima negotia non interrompere, animo proposueram, antequam certo fierent

B *Semel mihi imaginabar me videre figuram quandam, ad magnitudinem arenæ quam internæ cuidam corporis nostri parti comparare poteram. Cum materia hæc per momenta quedam aëri fuisset exposita, prædicta vasum multitudo in aquosam magnis oleaginosis globulis permixtam, materiam mutabatur: quales globulos inter medulla spinalis vasa interjacere antehac dixi. Hisce oleaginosis globulis visis mihi imaginabar, quod forsitan fuerint vasa convehendis spiritibus animalibus inservientia: eaque ex tam molli consistere materia, ut, intermittente humoris vel spirituum animalium transfluxu, illic in globulos oleaginosos diversa magnitudinis coalescant; præcipue cum aëri exponuntur. Et cum prædicta materia paucillum temporis steterat, in ea observabantur trilaterales figuræ ab utraque parte in aculeum desinentes, quibusdam longitudo minutissimæ arenæ, aliquæ aliquantum majores, ut fig. A. Præterea, adeo nitidæ ac pellucidæ, ac si crystallinæ fuissent.*



Figure 1: Parts of the original publication [1] from 1677 by Antoni van Leeuwenhoek with the description of the first light microscopic observation of crystalline spermine phosphate in human semen. (A) The title page 1040. (B) Page 1042 with fig. A showing the characteristic crystalline shape [2] of spermine $\times 2 H_3PO_4 \times 6 H_2O$ [2]. The last paragraph including fig. A is read in New Latin: “Et sum prædicta materia paucillum temporis steterat, in ea observabantur trilaterales figuræ ab utraque parte in aculeum desinentes, quibusdam longitudo minutissimæ arenæ, aliquæ aliquantum majores, ut fig. A. Præterea, adeo nitidæ ac pellucidæ, ac si crystallinæ fuissent.”. English transcription: “And I mentioned the matter which stood for a short time, in which trilateral figures were observed from both sides ending in a sting, some in length of minute grains, some a little larger, as fig. A. Moreover, so sleek and translucent, as if it were crystalline.”.

A

$$pK'_{a,HG} = \frac{8}{3} (n - 1) + pK_{a,R-OH} = 2.5601$$

$$\text{with } (n - 1) = \frac{0.821 \times 10^{-5} \text{ cm} \times \lambda}{\sqrt[3]{2 \lambda' s j b^2}}, \text{ and } \lambda = 1 + \ln \frac{3h_0^2}{2h_0^2}; \lambda' = \ln \frac{3h_0^2}{2h_0^2} - 1$$

B

$$pH_{DNA} = -\log_{10} \sqrt{10^{-pK'_{a,HG}} \times c_{DNA} \times 3,099,441,038 \times 2 \times 2.700\%} = 2.3208$$

$$pH_{spermine} = -\log_{10} \sqrt{\frac{10^{-14} \times 10^{-pK'_{spermine}}}{c_{DNA} \times \frac{1}{4} \times 3,099,441,038 \times 2.700\%}} = 10.2177$$

$$pH_{shell} = -\log_{10} \sqrt{\frac{10^{-14} \times 10^{-pK'_{shell}}}{c_{DNA} \times \frac{2}{3} \times 3,099,441,038 \times 2.700\%}} = 10.2019$$

Figure 2: The calculation of the apparent acid dissociation constant of the haploid human genome, and of the interphase intranuclear micro-pH [41] values induced by human genome depending on its intrinsic epigenetic occupation status. (A) The calculation of the apparent (effective) $pK'_{a,HG} = 2.5601$ ($\theta = 25^\circ C$) of the haploid human genome according to the method of *Katchalsky & Gillis* [30] and *Kuhn & Kuhn* [31] [$pK'_{a,HG}$ = apparent acid dissociation constant ($\theta = 25^\circ C$) of haploid *H. sapiens* genome B-DNA double helix; $pK_{a,R-OH} = 1.29$ = theoretical pK_a ($\theta = 25^\circ C$) [29] of one isolated internucleotide phosphodiester (R–OH) proton; $s = 1$ = number of statistical subunits on thread molecule [31]; $j = 6$ = number of spacing atoms (at least distance) in one dsDNA repeating unit [31]; $b = 0.334 \times 10^{-7} \text{ cm}$ ($0.334 \pm 0.01 \text{ nm}$ [32]) = length rise in cm of one B-DNA repeating unit (helix rise/bp) in solution; h_0 = end-to-end distance of dsDNA at half-neutralization; h_0 = end-to-end distance of dsDNA]. (B) The theoretical micro-pH [41] values surrounding *H. sapiens* haploid (and diploid) interphase euchromatin were calculated as the intranuclear micro-pH induced by human haploid (and diploid) genome treated as a weak acid, from pH_{DNA} , $pH_{spermine}$, and pH_{shell} , by applying the formula for pH induced by weak acids: $pH = -\log_{10} (K_a \times c_a)^{0.5}$, or the formula for pH induced by weak bases: $pH = -\log_{10} [(K_w \times K_b) \times (c_b)^{-1}]^{0.5}$ (K_a , acid dissociation constant; c_a , concentration weak acid; K_w , dissociation constant of water, $K_w = 10^{-14}$ ($\theta = 25^\circ C$); c_b , concentration weak base). The single-stranded human genome intranuclear interphase concentration c_{DNA} was calculated as: [1 genome $\times (6.022140857 \times 10^{23} \text{ mol}^{-1})^{-1}] \times (33.5268 \text{ fl})^{-1} = 49.5282 \text{ pM}$. The intranuclear micro-pH was calculated by the law for the calculation of the solution pH induced by salts of weak acids with weak bases: $pH_{salt} = 0.5 \times (pH_{acid} + pH_{base})$.

Calculation of the Hypothetical Intranuclear Micro-pH Mediated by Single Spermine Occupation of Human Interphase Euchromatin

The reference *H. sapiens* genome size was taken as 3,099,441,038 bp (RefSeq assembly accession number GCF_000001405.40) [12]. The number of base pairs for the number of protein-coding exons in *H. sapiens* genome (59,265) [12] was calculated with the median of amino acid residues/gene (470.6745 aa) (see Section 2.5): 470.6745 aa × 59,265 × 3 = 83,683,573 bp (2.7000% of *H. sapiens* genome). The intrinsic pH of *H. sapiens* haploid interphase genome was calculated with the volume of a human blood lymphocyte interphase nucleus taken as 33.5268 fl [femtoliter = (μm)³] [8, 9] after technical correction ($f = 0.834$) [8] of the published value (40.2 ± 2.0 fl) [9]. Because *H. sapiens* dsDNA haploid genome is to be treated as a weak acid [$pK'_{aHG} = 2.5601$ ($\vartheta = 25$ °C)] (Figure 2A), the hypothetical nuclear $pH_{DNA} = 2.3208$, mediated by *H. sapiens* haploid interphase genome dsDNA, can be calculated (Figure 2B). The concentration c_{DNA} was calculated as: [1 genome × (6.022140857 × 10²³ mol⁻¹)⁻¹] × (33.5268 fl)⁻¹ = 49.5282 pM. This concentration was multiplied by the number of phosphodiester moieties in dsDNA (3,099,441,038 × 2) and corrected for actively transcribed gene regions of *H. sapiens* genome (2.7000%). The four pK'_a values (± s.d.) of spermine $pK'_{a1} = 10.86 ± 0.06$, $pK'_{a2} = 10.05 ± 0.01$, $pK'_{a3} = 8.82 ± 0.01$, $pK'_{a4} = 7.95 ± 0.01$ [potentiometric titration in H₂O, $\vartheta = 25$ °C, ionic strength (NaCl) 0.1] [34] give the apparent (effective) $pK'_{spermine} = 9.42 ± 0.02$ (mean ± s.d.) ($\vartheta = 25$ °C). Because the hypothetical nuclear concentration of spermine tetracation covering B-DNA is one quarter of c_{DNA} , if we assume one spermine molecule covering four base pairs (single occupation) of B-DNA [35], and correcting for actively transcribed gene regions of *H. sapiens* genome (2.7000%), the $pH_{spermine} = 10.2177$ can be calculated (Figure 2B). Therefore, the hypothetical intranuclear micro-pH surrounding *H. sapiens* haploid interphase euchromatin when covered (single quartet occupation) by the spermine tetracation can be calculated as: $pH_{DNA/spermine} = 0.5 × (2.3208 + 10.2177) = 6.2693$. For the diploid dsDNA genome, after completed S phase during interphase, the hypothetical micro-pH surrounding *H. sapiens* diploid interphase euchromatin when covered (single quartet occupation) by the spermine tetracation is, again, $pH_{DNA/spermine} = 6.2693$.

Calculation of the Theoretical Intranuclear Micro-pH Mediated by Sperminium Hydrogen Phosphate/Cyclooctaoxygen Sodium Complex Occupation of Human Interphase Euchromatin

The apparent $pK'_{spermine}$ was corrected for the monohydrogen phosphate complexation to get the apparent (effective) pK'_{shell} of the sperminium hydrogen phosphate/

cyclooctaoxygen sodium complex epigenetic shell. Ortho-phosphoric acid H₃PO₄ shows the $pK'_{a1} = 2.161$, $pK'_{a2} = 7.207$, and $pK'_{a3} = 12.325$ ($\vartheta = 25$ °C) [36]. Therefore, the $pK'_{spermine}$ is to be corrected for the monohydrogen phosphate of the epigenetic shell: $pK'_{shell} = [9.42 + ((2.161 + 7.207) × 0.5) + 12.325] × 0.5 = 8.9623$. Because the theoretical nuclear concentration of the sperminium hydrogen phosphate/cyclooctaoxygen sodium complex covering dsDNA is two-third of c_{DNA} , when we assume one epigenetic complex molecule covering three base pairs on both strands of dsDNA (double occupation), and correcting for actively transcribed gene regions of *H. sapiens* genome (2.7000%), the $pH_{shell} = 10.2019$ (Figure 2B). Therefore, the theoretical intranuclear micro-pH surrounding *H. sapiens* haploid interphase euchromatin when covered (double triplet occupation) by the sperminium hydrogen phosphate/cyclooctaoxygen sodium complex can be calculated as: $pH_{DNA/shell} = 0.5 × (2.3208 + 10.2019) = 6.2614$. For the diploid dsDNA genome, after completed S phase during interphase, the theoretical micro-pH surrounding *H. sapiens* diploid interphase euchromatin when covered (double triplet occupation) by the sperminium hydrogen phosphate/cyclooctaoxygen sodium complex is, again, $pH_{DNA/shell} = 6.2614$.

Color Assay for Cyclo-O₈-Na⁺ Contained in RC – Destruction of Cyclo-O₈-Na⁺ by the Glyphosate Metabolite (Aminomethyl)Phosphonic Acid

Stock preparations were: KI (52 mg KI in 2,000 μl H₂O) (1), KI + starch (52 mg KI + 18 mg starch in 2,000 μl H₂O) (2), RC + KI (34 mg RC + 52 mg KI in 2,000 μl H₂O) (3), RC + KI + AMPA (34 mg RC + 52 mg KI + 22 mg AMPA in 2,000 μl H₂O) (4), RC + KI + starch (34 mg RC + 52 mg KI + 18 mg starch in 2,000 μl H₂O) (5), RC + KI + starch + AMPA (34 mg RC + 52 mg KI + 18 mg starch + 22 mg AMPA in 2,000 μl H₂O) (6). The concentrations in solution were: RC, 16.95 mM (with cyclo-O₈-Na⁺, 67.79 mM); KI, 156.63 mM; AMPA, 99.06 mM. The stock solutions were incubated at room temperature (RT, $\vartheta = 14.0$ °C) for 30 min, then at elevated temperature ($\vartheta = 25.2$ °C) for 2 min, after which time the first photograph series (Figure 3) was taken. The solutions were then incubated at elevated temperature for 48 min, after which time the second photograph (Figure 3) series was taken. The solutions were further incubated at RT for 14 h. Afterwards, both RC + KI (3) and RC + KI + AMPA (4) were mixed with 1,000 μl deuterated chloroform (CDCl₃), and extracted by shaking. Concomitantly, RC + KI + starch (5) and RC + KI + starch + AMPA (6) were both mixed with 8 mg of solid L-ascorbic acid (concentration of L-ascorbic acid in solution 22.71 mM). The solutions were succeedingly incubated at elevated temperature for 15 min, after which time the third photograph series (Figure 3) was taken.

Control Color Assay for Potential Reduction of Elemental Iodine by the Glyphosate Metabolite (Aminomethyl)Phosphonic Acid

Stock preparations were: blank without AMPA (1,000 μ l H₂O), AMPA stock solution (34 mg AMPA in 1,000 μ l H₂O), and two iodine stock solutions [each 80 mg elemental iodine in 2,000 μ l 90% (v/v) aqueous ethanol]. The stock preparations were incubated at room temperature (RT, $\vartheta = 14.0$ °C) for 15 min with occasional shaking, then at elevated temperature ($\vartheta = 25.2$ °C) for 15 min, until the stock preparations were nearly dissolved (AMPA full, iodine not fully). Then the AMPA stock solution was injected into the first iodine stock solution (it results solution **A**), no decoloration resulted. The H₂O blank was injected into the second iodine stock solution (it results solution **B**), no decoloration resulted. The concentrations in solution were now: AMPA, 102.07 mM; iodine as I₂, 105.07 mM. After incubation at elevated temperature for 15 min, into both solutions 1,000 μ l of water were injected, crystallization of elemental iodine followed, and no decoloration resulted. The first photograph series (Figure 3) was taken after 25 min incubation at elevated temperature. After incubation at elevated temperature for 15 min, 2,000 μ l of 90% (v/v) aqueous ethanol were injected in both **A** and **B**. The mixtures were shaken, the iodine dissolved to give clear deep brown solutions, and no decoloration resulted. After 5 min incubation at elevated temperature the second photograph (Figure 3) series was taken. Both **A**



Figure 3: Color assays for cyclo-O₈-Na⁺ contained in RC, for the destruction of cyclo-O₈-Na⁺ by the glyphosate metabolite (aminomethyl)phosphonic acid (AMPA) (left, 1–6), and for the potential reduction of elemental iodine by AMPA (right, A and B). Solutions (left, 1–6) were: KI (1), KI + starch (2), RC + KI (3), RC + KI + AMPA (4), RC + KI + starch (5), and RC + KI + starch + AMPA (6). The concentrations in solution were: RC, 16.95 mM (with cyclo-O₈-Na⁺, 67.79 mM); KI, 156.63 mM; AMPA, 99.06 mM. The solutions were incubated at two room temperatures for prolonged time. Afterwards (left, bottom row), both RC + KI (3) and RC + KI + AMPA (4) were extracted with deuterated chloroform (bottom phase), and (left, bottom row) both RC + KI + starch (5) and RC + KI + starch + AMPA (6) were treated with L-ascorbic acid. Legend: left, top row (1–6) = first photograph series; left, middle row (1–6) = second photograph series; left, bottom row (1–6) = third photograph series; right, top row (first photograph series): (A) AMPA (76.55 mM) + iodine (as I₂, 78.80 mM), (B) iodine (as I₂, 78.80 mM); right, bottom row (second photograph series): (A) AMPA (51.03 mM) + iodine (as I₂, 52.53 mM), (B) iodine (as I₂, 52.53 mM).

and **B** did not show any further change at RT during 24 h observation.

Color Assay for Cyclo-O₈-Na⁺ Contained in RC – Destruction of Cyclo-O₈-Na⁺ by Glyphosate and ROUNDUP®

Stock preparations were: RC + KI + starch (34 mg RC + 57 mg KI + 18 mg starch in 2,000 μ l H₂O) (1), RC + KI + starch + glyphosate (34 mg RC + 57 mg KI + 18 mg starch + 34 mg glyphosate in 2,000 μ l H₂O) (2), RC + KI + starch + glyphosate-Na (34 mg RC + 57 mg KI + 18 mg starch + 87 mg ROUNDUP® GRAN in 2,000 μ l H₂O) (3), RC + KI + glyphosate (34 mg RC + 57 mg KI + 34 mg glyphosate in 2,000 μ l H₂O) (4), RC + KI + glyphosate-Na (34 mg RC + 57 mg KI + 87 mg ROUNDUP® GRAN in 2,000 μ l H₂O) (5) [a saturated solution of 18 mg glyphosate (free acid) in 1,000 μ l of H₂O showed pH 1.5 ($\vartheta = 14.0$ °C)]. The concentrations in solution were: RC, 16.95 mM (with cyclo-O₈-Na⁺, 67.79 mM); KI, 171.69 mM; glyphosate, 100.55 mM; glyphosate-Na, 108.15 mM. The stock solutions were incubated at room temperature (RT, $\vartheta = 14.7$ °C) for 30 min, then at elevated temperature ($\vartheta = 25.2$ °C) for 70 min, after which time the photograph series (Figure 4) was taken. A control color assay for potential reduction of elemental iodine by glyphosate and ROUNDUP® GRAN was performed. Stock preparations were: blank without glyphosate [75 mg elemental iodine in 2,000 μ l 45% (v/v) aqueous ethanol], glyphosate stock solution [52 mg glyphosate in 2,000 μ l 45% (v/v) aqueous ethanol], and ROUNDUP® GRAN stock solution [124 mg ROUNDUP® GRAN in 2,000 μ l 45% (v/v) aqueous ethanol]. The stock preparations were incubated at room temperature (RT, $\vartheta = 14.1$ °C) for 45 min with occasional shaking. Then 75 mg solid elemental iodine were added to both the glyphosate and ROUNDUP® GRAN stock preparation. After incubation at elevated temperature for 1.5 h, the stock preparations were nearly dissolved (ROUNDUP® GRAN full, glyphosate and iodine not fully). Then 1,000 μ l of 90% (v/v) aqueous ethanol were injected into each preparation [resulting in solution **A1** (glyphosate), **A2** (glyphosate-Na), and **B** (blank)], no decoloration resulted. The concentrations in solution were now: glyphosate, 102.52 mM; glyphosate-Na, 102.77 mM; iodine as I₂, 98.50 mM. The photograph series (Figure 4) was taken after 6 h incubation at elevated temperature, no decoloration resulted. Both **A1**, **A2**, and **B** did not show any further change at RT during 24 h observation.

A color assay for cyclo-O₈-Na⁺ contained in NC was accordingly performed. The preparation was: NC + KI + starch (25 mg NC + 43 mg KI + 18 mg starch in 1,000 μ l H₂O). The concentrations in solution were: NC, 18.25 mM (with cyclo-O₈-Na⁺, 18.25 mM); KI, 259.04 mM. The preparation was incubated at elevated temperature for 10 h, after which time the first photograph (Figure 4) series was taken. The preparation was further incubated at room temperature (RT, ϑ

= 13.7 °C) for 40 h, after which time the second photograph series (Figure 4) was taken.

Enzyme Assay of the Glyphosate Metabolite (Aminomethyl)Phosphonic Acid with Human Mitochondrial γ -Aminobutyric Acid Transaminase

Recombinant human (16p13.2) mitochondrial 4-aminobutyrate aminotransferase (ABAT) (E.C. 2.6.1.19), mature full length protein aa 29–500 with N-terminal His-SUMO-tag and C-terminal Myc-tag, was purchased from MyBioSource, Inc. (San Diego, CA, USA). The sequence (472 aa) is: S Q E L M K Q L N I I Q N A E A V H F F C N Y E E S R G - N Y L V D V D G N R M L D L Y S Q I S S V P I G Y S H - P A L L K L I Q P Q N A S M F V N R P A L G I L P P E N F V E - K L R Q S L L S V A P K G M S Q L I T M A C G S C S N E N A L K T I F M - W Y R S K E R G Q R G F S Q E E L E T C M I N Q A P G C P D Y S I L S - F M G A F H G R T M G C L A T T H S K A I H K I D I P S F D W P I A P - F P R L K Y P L E E F V K E N Q Q E E A R C L E E V E D L I V K Y R K - K K K T V A G I I V E P I Q S E G G D N H A S D D F F R K L R D I A R K H - G C A F L V D E V Q T G G G C T G K F W A H E H W G L D D P A D - V M T F S K K M M T G G F F H K E E F R P N A P Y R I F N T W L G - D P S K N L L L A E V I N I I K R E D L L N N A A H A G K A L L T G L L - D L Q A R Y P Q F I S R V R G R G T F C S F D T P D D S I R N K L I L - I A R N K G V V L G G C G D K S I R F R P T L V F R D H H A H L F L - N I F S D I L A D F K (PLP-binding K_{329} active site in bold).

The 500 aa ABAT precursor protein (4-aminobutyrate aminotransferase, mitochondrial isoform 1 precursor, *H. sapiens*, NCBI Reference Sequence NP_000654.2) additionally bears the N-terminal peptide 1–28. The mature




Figure 4: Color assays for the destruction of cyclo- O_8-Na^+ contained in RC by glyphosate and ROUNDUP® GRAN (top, 1–5), for the potential reduction of elemental iodine by glyphosate and ROUNDUP® GRAN (bottom, A1, A2, and B), and for cyclo- O_8-Na^+ contained in NC (bottom, NC1 and NC2). Solutions (top, 1–5) were: RC + KI + starch (1), RC + KI + starch + glyphosate (free acid) (2), RC + KI + starch + ROUNDUP® GRAN (3), RC + KI + glyphosate (free acid) (4), and RC + KI + ROUNDUP® GRAN (5). The concentrations in solution were: RC, 16.95 mM (with cyclo- O_8-Na^+ , 67.79 mM); KI, 171.69 mM; glyphosate, 100.55 mM; glyphosate-Na, 108.15 mM. The solutions were incubated at two room temperatures for prolonged time. Legend (bottom): (A1) glyphosate (free acid, 102.52 mM) + iodine (as I_2 , 98.50 mM), (A2) ROUNDUP® GRAN (glyphosate-Na, 102.77 mM) + iodine (as I_2 , 98.50 mM), (B) iodine (as I_2 , 98.50 mM), (NC1, NC2) NC (18.25 mM, with cyclo- O_8-Na^+ , 18.25 mM) + KI (259.04 mM) after 10 h (NC1) and 50 h (NC2) incubation.

ABAT (aa 29–500, 53.27 kDa) is responsible for the catabolism of γ -aminobutyric acid (GABA), an important, mostly inhibitory neurotransmitter in the central nervous system, into succinic semialdehyde. The active enzyme is a homodimer of 53 kDa subunits, each condensed to one pyridoxal 5'-phosphate (PLP). The human ABAT deficiency phenotype includes psychomotor retardation, hypotonia, hyperreflexia, lethargy, refractory seizures, and electroencephalographic (EEG) abnormalities. To investigate whether AMPA is a substrate for human mitochondrial GABA transaminase, experiments were performed with commercially available recombinant human GABA transaminase according to the procedure of Schor *et al.* [37] with some modifications. Incubations with 500 nmol AMPA, in an assay volume of 120 μ l did not show activity of GABA transaminase towards AMPA, while the control assay using 500 nmol ^{15}N -GABA as substrate did result in the formation of the expected enzyme product. Subsequent inhibition experiments, with co-incubations of fixed amounts (500 nmol) of ^{15}N -GABA with increasing amounts of AMPA (0–2,000 nmol), revealed that AMPA did not act as an inhibitor of the GABA transaminase-catalysed reaction of ^{15}N -GABA to succinic semialdehyde. These combined results strongly suggest that AMPA is not a substrate for human GABA transaminase.

Enzyme Assay of the Glyphosate Metabolite (Aminomethyl)Phosphonic Acid with Human Wild-Type Alanine:Glyoxylate Aminotransferase

Recombinant human alanine:glyoxylate aminotransferase (AGT) was expressed in *E. coli* and purified as described [38]. The enzyme at 5 μ M concentration was incubated with 100 mM AMPA at $\theta = 25$ °C in 100 mM potassium phosphate buffer pH 7.4. At various times (1, 2, 5, 22 h), aliquots were withdrawn and the reaction was stopped by adding trichloroacetic acid 10% (v/v). The total amount of PLP and pyridoxamine 5'-phosphate (PMP) was determined by HPLC analysis as previously described (Figure 5A) [39]. The product of the half-transamination of AMPA by AGT is phosphonoformaldehyde which can be further oxidized to phosphonoformic acid (phosphonoformate, foscarnet) (Figure 5B).

Software

Molecular modeling was performed with ACD/Chem Sketch version 2022.1.0 with integrated ACD/3D Viewer (Advanced Chemistry Development, Inc., Toronto, Ontario, Canada) and processed with Mercury 2022.3.0 [The Cambridge Crystallographic Data Centre (CCDC), Cambridge, United Kingdom]. Additional molecular modeling was performed with ACD/Chem Sketch version 12.01 with integrated ACD/3D Viewer (Advanced Chemistry Development, Inc., Toronto, Ontario, Canada) and processed with Mercury 3.1 version 3.1.1 [The Cambridge Crystallographic Data Centre (CCDC), Cambridge, United Kingdom].

Photographing

Original photographs of the color reactions (Figure 3, Figure 4) were taken with the Kodak EasyShare M550 digital camera (Eastman Kodak Company, Rochester, NY, USA) and transferred to computer with the software Adobe PhotoDeluxe 2.0 Business Edition 1.0 (Adobe Systems Incorporation, San Jose, CA, USA).

Results

Calculation of the Genomic Coverage by the Cyclooctaoxygen Sodium-Bridged Spermine Phosphate Epigenetic Shell of Interphase DNA in Bovine Lymphocytes

The theoretical intracellular concentration of the sperminium hydrogen phosphate/cyclo- O_8 - Na^+ complex required to cover all triplets of the dsDNA genome in a blood lymphocyte of *Bos taurus* was calculated as 13.81 mM. The actual concentration of spermine was measured as 1.57 ± 0.12 (mM \pm s.d.) [11]. Therefore, the genomic dsDNA coverage of *B. taurus* genome can be calculated as 2.62 ± 0.50 (% \pm s.d.), since one unit of sperminium hydrogen phosphate/cyclo- O_8 - Na^+ complex is assumed to cover three nucleotides. A good correlation was obtained when this value was compared to the proportion of protein-coding exons in *B. taurus* genome which was calculated as 2.58%. For comparison, the human genome contains 2.70% protein-coding exons (see Section 2.5). This pointed to complete coverage of actively transcribed gene regions in *B. taurus* interphase genome by the sperminium hydrogen phosphate/cyclo- O_8 - Na^+ complex.




Figure 5: The catabolism of AMPA by human peroxisomal alanine:glyoxylate aminotransferase (AGT). (A) Time course of the AMPA half-transamination reaction of human AGT. The enzyme at a concentration of 5 μ M was incubated at $\theta = 25$ °C in 100 mM potassium phosphate buffer (pH 7.4). At the indicated times, aliquots were withdrawn and denatured. After removal of the precipitated protein by centrifugation, the supernatants were subjected to HPLC analysis (squares, PLP; circles, PMP). (B) The generation of phosphonoformate (foscarnet) from AMPA over the intermediate phosphonoformaldehyde by the rate-limiting transamination of the major environmental glyphosate metabolite AMPA. Phosphonoformaldehyde, seen as a glyoxylate analogue, could be oxidized by glyoxylate oxidase (GO) and lactate dehydrogenase (LDH) [66]. Foscarnet is a potent inhibitor of eukaryotic DNA polymerase α [67, 68], an enzyme crucially involved in maintaining chromosomal integrity and telomere length [69].





Figure 6: Molecular modeling of the postulated [5] first epigenetic shell of euchromatic *in vivo* DNA, as exemplified for a single-stranded hexanucleotide d(ApCpCpApTpGp), part of the transcribed and translated human gene open reading frame initiation codon ATG encrypted in the Kozak consensus sequence 5'-ACCATGG-3' [28], introducing a molecular biological model for sperminium hydrogen phosphate/cyclo- O_8 - Na^+ /ssDNA (left) and sperminium hydrogen selenite/cyclo- O_8 - Na^+ /ssDNA (right) interactions. Element color codings: grey, carbon; white, hydrogen; blue, nitrogen; red, oxygen; purple, phosphorus; green, sodium; yellow, selenium.

Calculation of the Spermine Coverage of Highly Condensed Mitotic Metaphase DNA in HeLa S3 cells

The content of spermine in HeLa S3 cell metaphase chromatin was calculated as 135.9 ± 16.1 pmol spermine/82.84 zmol dsDNA, and 116.1 ± 11.8 pmol spermidine/82.84 zmol dsDNA [15]. This corresponds to 1.64×10^9 molecules spermine *pro* one HeLa S3 cell dsDNA genome, and 1.40×10^9 molecules spermidine *pro* one HeLa S3 cell dsDNA genome. Since one spermine molecule is assumed to cover six base pairs (in the pure spermine form of A-DNA duplex [16] and Z-DNA duplex [17]), and one spermidine molecule is assumed to cover six base pairs (in the pure spermidine form of Z-DNA duplex [18]), this corresponds to a genomic coverage of 50.4% by the spermine tetracation, and of 43.0% by the spermidine trication. This accounts for 93.4% polyamine occupation of HeLa S3 cell dsDNA highly condensed metaphase chromatin by spermine and spermidine, calculated for six base pairs/polyamine unit. As one spermine molecule, in one special occasion, was found to cover four base pairs of a unique B-DNA [35], these values could be anticipated as being lower, since chromosomal DNA is predominantly in the B-DNA form.

Calculation of the Polyamine Coverage of Maximally Condensed Mitotic Late Anaphase/Early Telophase DNA in Murine Cryptal Enterocytes

The elemental phosphorus content [w (P) in mmol/kg dry weight] in female *Mus musculus* strain C3H/HeJ cryptal enterocytic mitotic (late anaphase/early telophase) chromatin was published [19]. The obtained *in vivo* value corresponds to a genomic dsDNA coverage (calculated for 6 bp/polyamine) of 102.1% (100% coverage is 5.88 bp/polyamine), and a nuclear RNA coverage (calculated for 6 nucleotides/polyamine) of 159.5%, by the spermine tetracation and spermidine trication (spermine/spermidine ratio 0.85). This strongly points to a function of polyamine occupation for nuclear RNA, assuming 100% coverage as 3.76 nucleotides/polyamine molecule. These results, both for dsDNA and nuclear RNA [hnRNA with pre-mRNAs, snRNA, snoRNA, RNase P, RNase MRP, various ncRNAs (lncRNA) and other nuclear RNAs], are a logic consequence of the maximal condensation grade peaking in late anaphase/early telophase mammalian chromatin [40].

Calculation of the Hydrogen Selenite (HSeO_3^-) Coverage of Human Euchromatin DNA Specifically at ATG Start Codon Sequences

A perfect correlation was obtained when the hydrogen selenite substitution content of the cyclooctaoxygen sodium-bridged sperminium hydrogen phosphate protection shell of human euchromatin was calculated from the optimal human blood serum level for the essential human micronutrient selenium (see Section 2.5). The optimal human blood serum level of selenium (as Se) was taken as 110 $\mu\text{g/l}$ (ng/ml) from literature [23, 24]. The median number of aa-coding triplets required for one triplet to be covered with the sperminium hydrogen selenite/cyclo- $\text{O}_8\text{-Na}^+$ complex in interphase euchromatin (in form of hydrogen selenite HSeO_3^-) at the published [23, 24] optimal human blood serum level of selenium [as Se, 110 $\mu\text{g/l}$ (ng/ml)] was calculated to be 560.44 aa. The optimal human blood serum level to cover all ATG triplet codons in human euchromatin mRNA genes, both at the mRNA gene translation initiation triplet ATG codon and at the 5'-UTR uORFs upstream start codons (uATGs transcribed to uAUGs), was calculated to be 107 $\mu\text{g/l}$ (ng/ml). The number of 5'-UTR uORFs upstream start codons (uATGs transcribed to uAUGs) in the 5'-UTR of mRNA genes was calculated from literature [26] as the number of transcribed human genes with at least one uATG divided by the number of human mRNA genes with annotated 5'-UTR. 49% of human mRNA genes contain at least one uATG start codon in their 5'-UTR [26] [the downstream 3'-untranslated region (3'-UTR) contains none ATG start codon]. The number of triplets required for the covering of all ATG triplet codons in human interphase euchromatin mRNA genes by one hydrogen selenite HSeO_3^- molecule (within the sperminium

hydrogen selenite/cyclo- $\text{O}_8\text{-Na}^+$ complex) at the calculated optimal human blood serum level of selenium 107 $\mu\text{g/l}$ (ng/ml) (576.40 aa) correlated with the determined heuristic value (560.44 aa). Consequently, a model of the postulated [5] first epigenetic shell of euchromatic *in vivo* DNA, as exemplified for a single-stranded hexanucleotide d(ApCpCpApTpGp), part of the transcribed and translated human gene open reading frame initiation codon ATG encrypted in the Kozak consensus sequence 5'-ACCATGG-3' [28], introducing a molecular biological model for sperminium hydrogen phosphate/cyclo- $\text{O}_8\text{-Na}^+$ /ssDNA and sperminium hydrogen selenite/cyclo- $\text{O}_8\text{-Na}^+$ /ssDNA interactions, was deduced (Figure 6).

Calculation of the Apparent Acid Dissociation Constant of the Human Genome DNA, and the Intranuclear Micro-pH Mediated by Single Spermine Occupation, in Comparison to Spermine Phosphate/Cyclooctaoxygen Sodium Complex Occupation, of Human Interphase Euchromatin

The apparent (effective) $\text{pK}'_{\text{a,HG}} = 2.56$ ($\theta = 25^\circ\text{C}$) of the haploid human genome was calculated (Figure 2A) according to the method of Katchalsky & Gillis [30], as based on the theoretical considerations of Kuhn & Kuhn [31]. The hypothetical intranuclear $\text{pH}_{\text{DNA}} = 2.32$, mediated by *H. sapiens* haploid interphase genome dsDNA without any neutralizing shell, can be calculated (Figure 2B). Assuming one spermine molecule covering four base pairs (single occupation) of B-DNA [35], and correcting for actively transcribed gene regions of *H. sapiens* genome, the hypothetical intranuclear micro-pH [41] surrounding *H. sapiens* haploid interphase euchromatin when covered (single quartet occupation) by the spermine tetracation alone can be calculated $\text{pH}_{\text{DNA/spermine}} = 6.27$. For the diploid dsDNA genome, after completed S phase during interphase, the $\text{pH}_{\text{DNA/spermine}}$ is identical. The theoretical intranuclear micro-pH surrounding *H. sapiens* haploid interphase euchromatin when covered (double triplet occupation) by the sperminium hydrogen phosphate/cyclooctaoxygen sodium complex can be calculated $\text{pH}_{\text{DNA/shell}} = 6.26$. For the diploid dsDNA genome, after completed S phase during interphase, the $\text{pH}_{\text{DNA/shell}}$ is identical.

Color Assay for Cyclo- $\text{O}_8\text{-Na}^+$ Contained in RC – Destruction Of Cyclo- $\text{O}_8\text{-Na}^+$ by the Glyphosate Metabolite (Aminomethyl)Phosphonic Acid

The cyclo- $\text{O}_8\text{-Na}^+$ complex [μ -chloro(μ -hydroxy) bis(octoxocane- $\kappa^4\text{O}^1, \text{O}^3, \text{O}^5, \text{O}^7$)disodium] contained in RC (4 mol cyclo- $\text{O}_8\text{-Na}^+$ *pro* mol cytidine \times HCl) reacted with potassium iodide and potato starch to an intensely colored (reddish violet) [(cyclo- $\text{O}_8\text{-Na}^+$) $_2(\text{I}_4^{2-})$]-amylose complex [RC + KI + starch (5)] (Figure 3) which was destroyed by the glyphosate metabolite (aminomethyl)phosphonic acid (AMPA) [RC + KI + starch + AMPA (6)]. The nature of this complex is based on the starch-catalysed formation of




Figure 7: Explanation of the color reaction for cyclo- O_8-Na^+ contained in **RC**, and the destruction of cyclo- O_8-Na^+ by the glyphosate metabolite (aminomethyl)phosphonic acid (AMPA). (A) The cyclo- O_8-Na^+ complex contained in **RC** reacted with potassium iodide under catalysis by potato starch to an intensely colored (reddish violet) $[(cyclo-O_8-Na^+)_2(I_4^{2-})]$ complex stabilized within the starch-contained amylose helix. (B) The proposed chemical formula for the amylose-complexed $[(cyclo-O_8-Na^+)_2(I_4^{2-})]$: bis(octoxocane- κ^4O^1,O^3,O^5,O^7)($\mu-2\lambda^3,3\lambda^3$ -tetraioda-1,3-diene- $\kappa^I^1:\kappa^I^2$)disodium. (C) A logically deduced catalytic 'rolling-circle' mechanism for the AMPA-catalysed degradation of cyclo- O_8-Na^+ . The cyclooctaoxygen ring is split to a phosphonate-esterified nonaoxidanide which is stabilized by ionic binding to the primary ammonium cation of AMPA. The phosphonate-esterified nonaoxidanide eliminates four oxygen O_2 molecules by a 'rolling-circle' cascade, regenerating AMPA.

tetraiodide I_4^{2-} $[(I-I-I-I)^{2-}]$, which is known to be of ruby red color in crystalline form [42], was frequently observed in crystals [43], and was theoretically predicted to exist in solution [44]. The tetraiodide I_4^{2-} is in turn complexed to cyclo- O_8-Na^+ and inserted into the amylose helix (Figure 7A and 7B). Multiple controls excluded that the $[(cyclo-O_8-Na^+)_2(I_4^{2-})]$ -amylose complex is formed (i) in blanks [KI (1), and KI + starch (2)], (ii) without starch [**RC** + KI (3)],

and (iii) from AMPA and **RC** + KI [**RC** + KI + AMPA (4)] (Figure 3). That the reddish violet complex involved triiodide I_3^- $[(I-I-I)^-]$ or pentaiodide I_5^- $[(I-I-I-I-I)^-]$ anions could be excluded by the observed color. Triiodide I_3^- is deep brown [45] and pentaiodide I_5^- is deep blue in color [46]. Pentaiodide I_5^- is also deep blue in the well-known complex with starch [47]. Furthermore, the $[(cyclo-O_8-Na^+)_2(I_4^{2-})]$ -amylose complex was (i) reduced (decolorized) by L-ascorbic acid (vitamin C), and (ii) the residual color after AMPA-catalysed destruction of cyclo- O_8 was pale pink (rosé) in color, not intense blue. This proved that (i) the reddish violet complex contained reducible iodine units, and (ii) did not contain the reducible iodine units as triiodide, pentaiodide and/or other higher polyiodides [48]. Without starch no $[(cyclo-O_8-Na^+)_2(I_4^{2-})]$ and/or iodine was formed, as was proved by deuterated chloroform ($CDCl_3$) extraction of the incubated **RC** + KI (3) and **RC** + KI + AMPA (4) solutions (Figure 3). Therefore, the iodine in tetraiodide I_4^{2-} must have being formed by starch catalysis.

A logically deduced catalytic 'rolling-circle' mechanism for the AMPA-catalysed degradation of cyclo- O_8 is hence proposed (Figure 7C). AMPA exhibits three acid dissociation constants: $pK_{a1} = 0.9$ (phosphonic acid, 1st), $pK_{a2} = 5.6$ (phosphonic acid, 2nd), $pK_{a3} = 10.2$ (primary ammonium $R-NH_3^+$) [49]. Therefore, AMPA is fully (zwitter)ionized at physiological pH 7.4. One anionic oxygen of the phosphonate group binds to the sodium cation in cyclo- O_8-Na^+ , the other anionic phosphonate oxygen splits the cyclooctaoxygen ring creating a phosphonate-esterified nonaoxidanide which is stabilized by ionic binding to the primary ammonium cation of AMPA. The phosphonate-esterified nonaoxidanide eliminates four oxygen O_2 molecules by a 'rolling-circle' cascade, regenerating AMPA.




Figure 8: A logically deduced catalytic 'rolling-circle' mechanism for the (fully ionized) glyphosate-catalysed degradation of cyclo- O_8-Na^+ . The cyclooctaoxygen ring is split to a phosphonate-esterified nonaoxidanide which is stabilized by ionic binding to the secondary ammonium cation of glyphosate (and complexation of the sodium cation). The phosphonate-esterified nonaoxidanide eliminates four oxygen O_2 molecules by a 'rolling-circle' cascade, regenerating glyphosate.

cascade, in reversal of the proposed [5] synthesis of cyclooctaoxygen, yielding AMPA and Na⁺. This would be clearly a catalytic mechanism, since AMPA is regenerated in the reaction cycle. Hence AMPA is able to destroy many cyclo-O₈-Na⁺ complexes without being consumed itself. To exclude that the destruction of cyclo-O₈-Na⁺ by AMPA is an artifact, it was tested if AMPA reduces (decolorizes) iodine in near equimolar mixture (Figure 3). In all variations tested, including a blank control, AMPA was not oxidized by iodine, and, in turn, did not reduce (decolorize) iodine molecules. Therefore, the AMPA-catalysed destruction of cyclo-O₈ was selective, and not a mere reduction of the tetraiodide I₄²⁻ [(I-I-I-I)²⁻]-contained iodine unit in the [(cyclo-O₈-Na⁺)₂(I₄²⁻)]-amylose complex.

Color Assay for Cyclo-O₈-Na⁺ Contained in RC – Destruction of Cyclo-O₈-Na⁺ by Glyphosate and ROUNDUP®

The developed color assay was applied onto the free acid of glyphosate and the monosodium salt of glyphosate contained in ROUNDUP® GRAN granules. The colored [(cyclo-O₈-Na⁺)₂(I₄²⁻)]-amylose complex [RC + KI + starch (1)] (Figure 4) was destroyed by glyphosate [RC + KI + starch + glyphosate (2)] and ROUNDUP® GRAN [RC + KI + starch + glyphosate-Na (3)]. Without starch no [(cyclo-O₈-Na⁺)₂(I₄²⁻)] and/or other color complex was formed, as




Figure 9: An improved and corrected model for the cyclooctaoxygen sodium-bridged spermine phosphate (and selenite) epigenetic shell [5] of actively transcribed gene regions in eukaryotic interphase 'open' chromatin DNA. (A) The nucleus of an eukaryotic cell with chromosome territories, chromatin fiber (10 nm 'beads-on-a-string' fiber), nucleosome structure, nucleosome octamer core histone proteins [H2A/H2B, H3/H3 variant, H4] with posttranslational histone protein modifications (histone PTM), and decondensing DNA with regulative cytosine nucleobase 5-methylation sites (me). Adapted and modified in part from [70]. (B) The decondensation of chromatin enabling gene transcription in eukaryotic interphase 'open' chromatin DNA. The relaxed chromatin, regulatory transcription factor, chromatin-remodeling complex/histone acetyltransferases (HATs), and chromatin remodeling are indicated. The 'open' DNA with the transcription-prone gene is generally structured in promoter, promoter-proximal genetic elements, transcribed/expressed gene exons and non-expressed gene introns. The cyclooctaoxygen sodium-bridged spermine phosphate (and selenite) epigenetic coverage of this actively transcribed gene region is symbolized.

was proved by the controls RC + KI + glyphosate (4) and RC + KI + glyphosate-Na (5) (Figure 4). The RC + KI + starch + glyphosate (2) solution was colored yellow by the strong acid glyphosate (Figure 4), through H⁺ action onto starch producing the characteristic yellow dextrans [50]. Glyphosate exhibits four acid dissociation constants: pK_{a1} = 0.78 (phosphonic acid, 1st), pK_{a2} = 2.29 (carboxylic acid), pK_{a3} = 5.96 (phosphonic acid, 2nd), pK_{a4} = 10.98 (secondary ammonium R₁-NH₂⁺-R₂) [49]. Therefore, glyphosate (free acid) represents a strong acid. A logically deduced catalytic 'rolling-circle' mechanism for the glyphosate-catalysed degradation of cyclo-O₈ is hence proposed (Figure 8). One anionic oxygen of the phosphonate group binds to the sodium cation in cyclo-O₈-Na⁺, the other anionic phosphonate oxygen splits the cyclooctaoxygen ring creating a phosphonate-esterified nonaoxidanide which is stabilized by ionic binding to the secondary ammonium cation of glyphosate. The phosphonate-esterified nonaoxidanide eliminates four oxygen O₂ molecules by a 'rolling-circle' cascade, yielding glyphosate and Na⁺ in a catalytic mechanism. Glyphosate is regenerated in the catalytic cycle. Glyphosate is able to destroy many cyclo-O₈-Na⁺ complexes without being consumed itself (Figure 8).

To exclude that the destruction of cyclo-O₈-Na⁺ by glyphosate is an artifact, it was tested if glyphosate, or glyphosate-Na, reduces (decolorizes) iodine in near equimolar mixture (Figure 4). Glyphosate was not oxidized by iodine, and, in turn, did not reduce (decolorize) iodine molecules. Therefore, the glyphosate-catalysed destruction of cyclo-O₈ was selective. To confirm the general nature of the color assay, it was extended to the cyclo-O₈-Na⁺ contained in NC (1 mol cyclo-O₈-Na⁺ *pro* 4 mol cytidine × HCl) [5]. The reddish violet [(cyclo-O₈-Na⁺)₂(I₄²⁻)]-amylose complex was indeed formed from NC (Figure 4), but much more slowly (> 10 h) than from RC.

Enzymatic Investigations with the Glyphosate Metabolite (Aminomethyl)Phosphonic Acid

AMPA was tested for being accepted as an enzymatic substrate for human mitochondrial γ-aminobutyric acid transaminase (ABAT) [37], and wild-type human liver peroxisomal alanine:glyoxylate aminotransferase (AGT) [38]. Both enzymes were selected because of the chemical analogy between AMPA and β-alanine/L-alanine. ABAT represents also a β-alanine transaminase [37]. ABAT did not catalyse any AMPA half-transamination. Data obtained with human AGT indicate that the enzyme is slowly able to catalyse the half-transamination of AMPA (Figure 5B), with a rate of $k_{cat} = -0.0108 \pm 0.0009 \mu\text{M PLP/h}/\mu\text{M AGT}$ consumed, or $k_{cat} = -0.0104 \pm 0.0009 \mu\text{M PMP/h}/\mu\text{M AGT}$ formed, respectively (PLP, pyridoxal 5'-phosphate; PMP, pyridoxamine 5'-phosphate) (Figure 5A). This catalytic constant is approximately 16.2 million-fold lower than that of

Table 1: Tabulation of the selective cell cycle-dependent occupation of eukaryotic DNA by epigenetic polyamine shells. 1 ×, single quartet occupation (one polyamine *pro* both strands); 2 ×, double triplet occupation (one polyamine *pro* one strand); Chr, chromatin.

Cell cycle phase		Heterochromatin		Euchromatin	
Interphase	Function (concise)	Spermine-occupation?	Cyclo-O ₈ -Na ⁺ -occupation?	Spermine-occupation?	Cyclo-O ₈ -Na ⁺ -occupation?
G ₀ (Gap 0)	Resting and quiescence	No	No	Yes (2 ×) – With cyclo-O ₈ -Na ⁺	Yes (2 ×) – On 'open' Chr
G ₁ (Gap 1)	Transcription and histone synthesis	No	No	Yes (2 ×) – With cyclo-O ₈ -Na ⁺	Yes (2 ×) – On 'open' Chr
S (Synthesis)	DNA synthesis	No	No	Yes (2 ×) – With cyclo-O ₈ -Na ⁺	Yes (2 ×) – On 'open' Chr
G ₂ (Gap 2)	Translation	No	No	Yes (2 ×) – With cyclo-O ₈ -Na ⁺	Yes (2 ×) – On 'open' Chr
Mitosis	Function (concise)	Spermine-occupation?	Cyclo-O ₈ -Na ⁺ -occupation?	Spermine-occupation?	Cyclo-O ₈ -Na ⁺ -occupation?
Prophase	Chr condenses into chromosomes, nucleolus disappears	Yes (1 ×) – Condensing Chr	No	Yes (1 ×) – Condensing Chr	No
Prometaphase	Kinetochores and polar microtubules attach, mitotic spindle formed, nucleus disappears	Yes (1 ×) – Condensed Chr	No	Yes (1 ×) – Condensed Chr	No
Metaphase	Centrosomes pull chromosomes, chromosome centromeres line up at metaphase plate	Yes (1 ×) – Highly condensed Chr	No	Yes (1 ×) – Highly condensed Chr	No
Anaphase	Chromosomes break at centromeres, sister chromatids separated by microtubules	Yes (1 ×) – Maximally condensed Chr in late anaphase	No	Yes (1 ×) – Maximally condensed Chr in late anaphase	No
Telophase	Chr reformed from chromosomes, nucleus and nucleolus reappear	Yes (1 ×) – Maximally condensed Chr in early telophase	No	Yes (1 ×) – Maximally condensed Chr in early telophase	No

the physiological transamination of L-alanine (human wild-type AGT: $k_{cat} = 45 \pm 2 \text{ s}^{-1}$; $k_{cat} = 162,000 \text{ h}^{-1}$) [51].

Discussion

One reason why the cyclooctaoxygen sodium-bridged spermine phosphate epigenetic shell of *in vivo* DNA was overlooked until now should be its destruction during DNA purification by the classical phenol extraction method of

Schuster, Schramm & Zillig [52]. This original phenol extraction, although variously modified for nowadays use, consistently precipitates the spermine as sperminediium (terminal NH₃⁺) di(phenolate) at pH 7.9–8.0, since the pK_a value of phenol is 9.97 [potentiometric titration in H₂O, $\theta = 25 \text{ }^\circ\text{C}$, ionic strength (NaCl) 0.1], and the four pK_a values of spermine are: pK_{a1} = 10.86 (terminal [53] NH₃⁺), pK_{a2} = 10.05 (terminal NH₃⁺), pK_{a3} = 8.82 (inner [53] NH₂⁺), pK_{a4} = 7.95

(inner NH_2^+) [potentiometric titration in H_2O , $\theta = 25^\circ\text{C}$, ionic strength (NaCl) 0.1] [34]. The cyclo- $\text{O}_8\text{-Na}^+$ could react with alkaline (pH 7.9–8.0) buffered phenol to disodium rhodizonate ($\text{C}_6\text{Na}_2\text{O}_6$), a known oxidation product of *p*-benzoquinone, which in turn is an oxidation product of phenol. Regardless of the chemical details, the commercial salmon sperm DNA (utilized in [5]) and calf thymus DNA preparations are devoid of cyclo- $\text{O}_8\text{-Na}^+$ and spermine phosphate complexation, since the methods utilized for calf thymus and salmon sperm sodium deoxyribonucleate preparation [treatment with sodium dodecyl sulfate, high salt (NaCl) treatment, repeated ethanol precipitation] certainly remove the cyclooctaoxygen sodium-bridged spermine phosphate epigenetic shell. Our findings have important consequences for the epigenetics [54] of eukaryotic *in vivo* DNA. Kesel *et al.* suggested [5] a model for a first epigenetic shell of *in vivo* DNA (Figure 6), based on the observed complexation of cyclo- $\text{O}_8\text{-Na}^+$ and spermine phosphate to ssDNA. In this model [5] the phosphate backbone of ssDNA binds one cyclo- $\text{O}_8\text{-Na}^+$ *pro* three nucleotides, and this binary complex binds one spermine monophosphate to form a ternary epigenetic core of DNA. The monohydrogen phosphate bridges the cyclo- $\text{O}_8\text{-Na}^+$ with the sperminium cation, and the cyclo- $\text{O}_8\text{-Na}^+$ has an inverted alternating orientation [5]. Interestingly, the sperminium tetracation cannot bind alone to DNA in this model, since the distances [$d(N^1, N^4) = 490$ pm; $d(N^4, N^9) = 620$ pm; $d(N^1, N^{12}) = 1,600$ pm] between the four ammonium nitrogens do not fit the intrastrand phosphate–phosphate distance of dsDNA (B-DNA: $d_o = 700$ pm [55]; A-DNA: $d_o = 590\text{--}600$ pm [55]; Z-DNA: $d_o = 590$ pm (step pCp), $d_o = 600$ pm (step pGp) [56]). Therefore, it is quite remarkable that in this model for the first epigenetic shell of *in vivo* DNA [5] a repeating unit is formed from cyclo- $\text{O}_8\text{-Na}^+$ and spermine phosphate that perfectly fits both the triplet nature of the genetic code and the repeating distance of the phosphate anion backbone of DNA. In view of the important findings of Kirmes *et al.* [4], that an interaction of eukaryotic chromatin DNA structure with atmospheric oxygen partial pressure takes place, previous postulations [5] have to be defined more precisely. Since under switching to hypoxic conditions eukaryotic cell chromatin gets highly condensed [4], accompanied by redistribution of the polyamine pool to the nucleus [4], the cyclooctaoxygen sodium-bridged spermine phosphate epigenetic shell can only be restricted to actively transcribed gene regions of eukaryotic ‘open’ euchromatin, excluding occupation of condensed chromatin. Hypoxia should largely prevent metabolic formation of cyclooctaoxygen. Both under hypoxic conditions and in the metaphase of mitosis, where spermine synthesis is highest [57], coincident with an extraordinary high condensation grade (15,000–20,000-fold) of metaphase chromatin [58], none cyclooctaoxygen should be involved in covering the highly condensed chromatin DNA. Here no discrimination between eu- and

heterochromatin is made in metaphase, and all mitotic eukaryotic chromatin DNA is complexed with spermine tetracation and spermidine trication (and, at small proportions, with putrescine and cadaverine dications). In summary, this reflects the high mitotic chromatin condensation grade and is confirming the results with hypoxia-induced chromatin condensation under coinciding polyamine pool nuclear translocation [4]. All these considerations point to the correctness of the model that the cyclooctaoxygen sodium-bridged spermine phosphate epigenetic shell is restricted to actively transcribed ‘hot spot’ gene regions of eukaryotic ‘open’ euchromatin. Importantly, this epigenetic shell of eukaryotic ‘open’ euchromatin covers each strand of dsDNA separately, one at the positive strand, one at the negative strand (double occupation), whereas the highly condensed dsDNA structures bind one polyamine molecule directly at the double strand (single occupation) [16–18, 35]. This is substantiated by the precise calculation of the apparent acid dissociation constant of the human genome DNA. Kesel *et al.* also elaborated a model for selenium (as hydrogen selenite, HSeO_3^- , at physiological pH 7.4) protection of DNA [5]. Selenium, the element of the moon [59], was discovered by Jöns Jacob Berzelius (1779–1848) in 1817 and was named by him in honor of the Greek goddess of the moon *Selene* (σελήνη) [60]. Selenium is essential to mammalian physiology at nutritional levels, but suprphysiological intake of selenium is known to be toxic for mammals [5]. Sodium selenite (Na_2SeO_3), as hydrogen selenite HSeO_3^- at pH 7.0 (selenious acid H_2SeO_3 ; $\text{p}K_{a1} = 2.62$, $\text{p}K_{a2} = 8.32$ [61]), binds to calf thymus genomic B-DNA at pH 7.0 [62], and to *Saccharomyces cerevisiae* A-RNA at pH 7.0 [63]. Selenium has the ability to protect DNA from noxious influences (oxidative stress, radiation, cytotoxic agents) [5], and is essential to genomic stability [5], but the exact molecular biological basis for these phenomena is unknown. If in the model of a first epigenetic shell of *in vivo* DNA [5] the monohydrogen phosphate is replaced by hydrogen selenite, an epigenetic explanation for the interaction of selenium with eukaryotic *in vivo* DNA could be given. A coinciding correlation was obtained when the hydrogen selenite substitution content of the cyclooctaoxygen sodium-bridged sperminium hydrogen phosphate protection shell of human euchromatin was calculated from the optimal human blood serum level for the essential human micronutrient selenium (see Section 2.5). The optimal human blood serum level of selenium (as Se), given as 110 $\mu\text{g/l}$ (ng/ml) from literature [23, 24], coincided with the calculated optimal selenium human blood serum level, 107 $\mu\text{g/l}$ (ng/ml), required to cover all ATG triplet codons in human euchromatin mRNA genes, both at the mRNA gene translation initiation triplet ATG codon and the 5'-UTR uORFs upstream start codons (uATGs transcribed to uAUGs). 49% of human mRNA genes contain at least one uATG start codon in their 5'-UTR [26] [the downstream 3'-untranslated region (3'-UTR) contains none ATG start

codon]. The median number of aa-coding triplets required for one triplet to be covered with the sperminium hydrogen selenite/cyclo- O_8 - Na^+ complex in interphase euchromatin (in form of hydrogen selenite $HSeO_3^-$) at the published [23, 24] optimal human blood serum level of selenium [as Se, 110 $\mu\text{g/l}$ (ng/ml)] was calculated as to be 560.44 aa. This result correlated well with the calculated number of aa triplets (576.40) required for the covering of all ATG triplet codons in human interphase euchromatin mRNA genes by one hydrogen selenite $HSeO_3^-$ molecule (within the sperminium hydrogen selenite/cyclo- O_8 - Na^+ complex) at the calculated optimal human blood serum level of selenium [as Se, 107 $\mu\text{g/l}$ (ng/ml)]. A model of the postulated [5] first epigenetic shell of euchromatic *in vivo* DNA was deduced, as exemplified for a single-stranded hexanucleotide d(ApCpCpApTpGp), part of the transcribed and translated human gene open reading frame initiation codon ATG encoded in the *Kozak* consensus sequence [28] (Figure 6). This model accounts for, at least some of, the well-known bimodal, protective and toxic, *in vivo* effects exerted by selenium onto mammalian physiology. A complete specific substitution pattern of hydrogen selenite for monohydrogen phosphate at ATG open reading frame initiation codons would be essential for eukaryotic gene regulation, but if the displacement ratio $HSeO_3^-/HPO_4^{2-}$ exceeds a certain physiological tolerance level, and hydrogen selenite is randomly incorporated into the eukaryotic protection shell of human euchromatin, the epigenetic equilibrium should collapse. The high, both acute and chronic, mammalian toxicity of sodium selenite (Na_2SeO_3) should be due, at least in part, to direct detrimental effects of supraphysiological levels of hydrogen selenite $HSeO_3^-$ on mammalian chromosomal DNA integrity and regulation of genome expression. Assuming an essential biological function for the cyclooctaoxygen sodium-bridged spermine phosphate and selenite epigenetic shell, it was searched for substances able to *selectively* destroy this epigenetic protection structure, and the total herbicide glyphosate, *N*-(phosphonomethyl)glycine (ROUNDUP®, Monsanto), and its major environmental metabolite (aminomethyl)phosphonic acid (AMPA) [64] were tested on the cyclo- O_8 - Na^+ complex contained in **RC**. Glyphosate was chosen because it represents the top-selling total herbicide worldwide [65], and **RC** was selected because of its highest molar cyclo- O_8 - Na^+ content in the complex series **NC**, **dNC**, and **RC** [5]. Glyphosate and AMPA show chemical properties which might predispose them for destruction of cyclooctaoxygen in general. Glyphosate and AMPA are very hydrophilic and amphoteric, and their phosphonate moieties could be suitable to interact with cyclo- O_8 - Na^+ . It was unequivocally shown that glyphosate and AMPA indeed *selectively* destroy the cyclo- O_8 - Na^+ complex contained in **RC** (Figure 3, Figure 4, Figure 7). I therefore conclude that glyphosate and the major environmental glyphosate metabolite AMPA [64] also destroy the cyclooctaoxygen

sodium-bridged spermine phosphate and selenite epigenetic shell of human euchromatin, because destruction of cyclooctaoxygen is sufficient to bring this essential protection shield of human euchromatin into collateral epigenetic collapse. To get support for the selectivity of AMPA as an epigenetic poison, the affinity of AMPA towards human mitochondrial γ -aminobutyric acid transaminase (ABAT) [37], and to wild-type human liver peroxisomal alanine:glyoxylate aminotransferase (AGT) [51], was determined. ABAT represents also a β -alanine transaminase [37], and both enzymes were selected because of the structural similarity between AMPA and β -alanine/L-alanine. AMPA showed essentially no affinity to ABAT, but was very slowly catabolized by AGT (Figure 5A). This latter result is of interest, since the product of the half-transamination of AMPA by AGT is phosphonoformaldehyde (Figure 5B) which can be oxidized (peroxisomal glycolate oxidase, cytoplasmic lactate dehydrogenase [66]) to phosphonoformic acid (phosphonoformate, foscarnet) (Figure 5B). Foscarnet represents a well-known inhibitor of mammalian [67, 68] and viral [67, 68] DNA-dependent DNA polymerases. Eukaryotic DNA polymerase α is crucially involved in chromosome maintenance, DNA repair and recombination, transcriptional silencing, checkpoint activation, and telomere length maintenance [69]. Mammalian DNA polymerase α is potently inhibited by foscarnet [67, 68]. Therefore, the low-affinity half-transamination of AMPA by AGT, the rate-limiting step leading to foscarnet, could negatively influence human chromosome maintenance, DNA damage repair, and telomere length preservation, mediated by the AMPA catabolite foscarnet inhibition of DNA polymerase α . This enzymatic catabolism provides an additional, minor mechanism of destabilization and impairment of eukaryotic chromosomal DNA indirectly induced by the environmental glyphosate metabolite AMPA.

Conclusions

An improved and corrected molecular biological model is proposed for a first epigenetic shell of eukaryotic euchromatin (Figure 6, Figure 9). This model incorporates an epigenetic explanation for the interactions of the essential micronutrient selenium (as selenite) with eukaryotic euchromatin (Figure 6). The sperminium hydrogen phosphate/cyclooctaoxygen sodium complex was calculated to cover the actively transcribed regions (2.6%) of bovine lymphocyte interphase genome dsDNA (double occupation). The polyamine (spermine/spermidine ratio 1.17) coverage of HeLa S3 cell metaphase chromatin dsDNA was calculated as 93.4% (single occupation). In murine cryptal enterocytic mitotic (late anaphase/early telophase) chromatin the obtained *in vivo* value corresponds to complete genomic coverage (single occupation), and to comprehensive and extensive nuclear RNA coverage, by the spermine tetracation and spermidine trication (spermine/spermidine ratio 0.85).




Figure 10: A chemical mechanistic deduced logical scheme for the generation of kinetin (*N*⁶-furfuryl-9*H*-adenine) [74] from DNA by proton catalysis (kinetin-generating “base flip”, KGBF). Protonation of the adenine nucleobase of the shown d(pApApAp) sequence at purine N-7 induces depurination [73], creating an apurinic site [73]. Proton-catalysed *N*-glycosylation [77, 78] at the 6-NH₂ of adenine regenerates a nucleobase-inverted 2'-deoxynucleotide which is in furanose ring-opening equilibrium with its ald(ox)imine form. Two eliminations simultaneously follow, the first (auto-catalysed by 3'-*O*-phosphate conjugate base proton reception) by S_N2-nucleophilic eliminative epoxide formation, the second by proton-catalysed (auto-catalysed by the previously liberated 3'-*O*-monohydrogen phosphate conjugate acid) β-elimination (E1 unimolecular elimination), which leads to an 3,4-epoxyolefin: *N*-{(1*E*,2*E*)-3-[(2*S*)-oxiran-2-yl]prop-2-en-1-ylidene}-9*H*-purin-6-amine. The chiral (*S*)-2-vinyloxirane structure rearranges to a chiral (*R*)-2-alkyl-2,5-dihydrofuran due to 3,4-epoxyolefin rearrangement [79, 80]: *N*-[(2*R*)-2,5-dihydrofuran-2-ylmethyl]-9*H*-purin-6-amine. The latter compound rearranges to kinetin through facile double bond migration driven by aromatization to a furan. A DNA single-strand break is left back.

Because cyclooctaoxygen seems to be naturally absent in hypoxia-induced highly condensed chromatin [4], a model [70] is proposed for the cyclooctaoxygen sodium-bridged spermine phosphate (and selenite) epigenetic shell of actively transcribed gene regions in eukaryotic ‘open’ chromatin DNA (Figure 9). Furthermore, a working model is tabulated in summary for the selective cell cycle-dependent epigenetic occupation of eukaryotic DNA (Table 1).

What may be the evolutionary conserved biological

significance, and human pathophysiological implication, of this selective epigenetic shell? Obviously, the selective cyclooctaoxygen sodium-bridged spermine phosphate (and selenite) epigenetic shell serves to discriminate euchromatin from heterochromatin (Table 1). Interestingly, a supraphysiological concentration of selenite induces S-G₂/M cell cycle arrest, locked in interphase [71], and inhibits DNA and RNA synthesis [72], again pointing on a crucial determination of the optimally hydrogen selenite-marked ATG codons in eukaryotic DNA mRNA genes.

Both a deficiency (not all ATG codons marked by HSeO_3^-) and an excess (randomly distribution of HSeO_3^- -tagging in genome, no more restricted to only ATG codons) of selenium concentration as hydrogen selenite exerts detrimental effects on eukaryotic gene regulation and expression. This reflects the specific life-essential role of selenium in eukaryotic gene processing and maintenance, a long-sought function of selenium in human nutritional physiology, previously defined as ‘mysterious’ essentiality of selenium for human life [59, 71].

Furthermore, during transcription of actively transcribed gene regions in eukaryotic ‘open’ chromatin the double helix must be unwound by DNA helicases and the strands must be separated to enable access to DNA-dependent RNA polymerases I, II, and III. This creates intermediate DNA single-strand regions which are prone to chemical structure damage by multiple noxious impacts like reactive oxygen species (ROS), mutagens, and UV light. The selective cyclooctaoxygen sodium-bridged spermine phosphate (and selenite) epigenetic occupation of these sensitive single-stranded stretches could serve as an intrinsic protection against structural damage. This would be a logic explanation for the selective nature of the separate occupation of both DNA strands, consequently retained when strands are separated for transcription of mRNA. Also intriguing seem to be pH effects, since spermine is a strong base, and the major pH-related damage to DNA is depurination [73] creating apurinic sites at low pH [73]. The precisely calculated intranuclear micro-pH gain, obtained by sperminium hydrogen phosphate/cyclooctaoxygen sodium complexation of B-DNA individual strands (essentially the same as the intranuclear micro-pH gain for condensed B-DNA strand-overarching covered by sperminium tetracations) should protect against low pH-depurination creating apurinic DNA lesions leading to DNA single-strand breaks [73]. In this context the formation of kinetin (N^6 -furfuryl-9H-adenine) from DNA is well known (Figure 10) [74]. It should be emphasized that kinetin is not contained in native mammalian DNA, contrary to misleading claims [75, 76], but is formed only during non-oxidative DNA damage. A mechanism for the kinetin formation in, or from, DNA was proposed [75, 76], but it seems not to be conclusive in chemical reason, since furfural does not react with the adenine 6-NH₂ group under condensation to a Schiff base under *in vivo* conditions [74].

I therefore propose a chemical mechanistic deduced logical scheme for the generation of kinetin from DNA by proton catalysis (kinetin-generating “base flip”, KGBF) (Figure 10), based on proton-catalysed depurination [73] and subsequent inverted adenine 6-NH₂ *N*-glycosylation [77, 78], in consequence [79, 80] leaving back a DNA single-strand break. It is therefore concluded that the sperminium hydrogen phosphate/cyclooctaoxygen sodium complex serves to protect ssDNA from nucleic acid-mediated intrinsic

low intranuclear micro-pH-induced depurination, including KGBF, creating apurinic sites and concomitant DNA single-strand breaks at eukaryotic genome regions engaged in active transcription. In conclusion, it is logically obvious that any chemical agent, biochemical precursor (selenium) deficiency, and/or physical circumstance compromising the sperminium hydrogen phosphate/selenite-cyclooctaoxygen sodium complexation will inevitably lead to a severe disturbance of eukaryotic genome integrity, to an increased mutation rate, and to genomic DNA single-strand breaks. This is, in part, proved by the *Snyder–Robinson* X-linked mental retardation syndrome [5, 81], characterized by a defect in spermine synthesis, leading to nearly complete loss of the polyamine spermine. Therefore, it was initiated a search for chemical agents selectively destroying the epigenetic shell of eukaryotic euchromatin. A candidate molecule was found, and, hence, it was defined as an ‘epigenetic poison’. The total herbicide glyphosate, *N*-(phosphonomethyl)glycine (ROUNDUP®, Monsanto), and its major environmental metabolite (aminomethyl)phosphonic acid (AMPA) [64] were found, rather unequivocally, to selectively destroy the cyclo-O₈-Na⁺ complex contained in RC. Glyphosate and AMPA came into focus because (i) glyphosate represents the top-selling total herbicide worldwide [65], (ii) their chemical structure (phosphonate + amine) and properties (strongly hydrophilic and acidic) seemed to enable them to interact with cyclooctaoxygen sodium, (iii) glyphosate and ROUNDUP® are suspected to damage DNA and cause cancer in humans [82], and (iv) AMPA is already widely distributed in global ecosystems like (surface) water [83]. I allow me to conclude on basis of my findings that glyphosate, ROUNDUP® and AMPA are major examples of slow-acting, insidious ‘epigenetic poisons’, (i) slowly eroding and deteriorating human, animal and plant genomic integrity, (ii) rattening human, animal and plant inborne protection of hereditary information against mutation, and (iii) disturbing the processing of human, animal and plant genetic information by transcription. It is hence inevitable for me to define glyphosate, ROUNDUP® and AMPA as a significant threat for human, animal and plant genomic stability, especially for future human generations forced to live under the glyphosate-, ROUNDUP®- and AMPA-induced radiomimetic effects.

Acknowledgments

The author is deeply indebted to T. Westfeld, E.-M. May, D. Wiegel, W. Wübbolt, O. Meier, R. Sachs, A. Karbach, W. Bergmeier, J. Moldenhauer and H.-J. Hühn (Currenta GmbH & Co. OHG, Leverkusen, Germany) for analytical services. I heartily thank H.J. Jodl for helpful discussions, E.A. Struys (University Hospital Vrije Universiteit, Amsterdam, The Netherlands) for performing the ABAT assays, and B. Cellini (University of Perugia, Department of Experimental Medicine, Italy) for conducting the AGT assays. I am obliged

to K. Hecker, K. Meuser and K. Hecker (HEKAtech GmbH, Wegberg, Germany) for expert elemental analyses.

References

- van Leeuwenhoek A. Observationes D. Anthonii Lewenhoeck, de natis è semine genitali animalculis. *Philos Trans R Soc Lond* 12 (1677–1678): 1040-1046.
- Rosenheim O. The isolation of spermine phosphate from semen and testis. *Biochem J* 18 (1924): 1253-1262.
- Cooke MS, Evans MD, Dizdaroglu M, et al. Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J* 17 (2003): 1195-1214.
- Kirmes I, Szczurek A, Prakash K, et al. A transient ischemic environment induces reversible compaction of chromatin. *Genome Biol* 16 (2015): 246.
- Kesel AJ, Day CW, Montero CM, et al. A new oxygen modification cyclooctaoxygen binds to nucleic acids as sodium crown complex. *Biochim Biophys Acta* 1860 (2016): 785-794.
- Igarashi K, Kashiwagi K. Polyamines: mysterious modulators of cellular functions. *Biochem Biophys Res Commun* 271 (2000): 559-564.
- Segel GB, Cokelet GR, Lichtman MA. The measurement of lymphocyte volume: importance of reference particle deformability and counting solution tonicity. *Blood* 57 (1981): 894-899.
- Kuse R, Schuster S, Schübbe H, et al. Blood lymphocyte volumes and diameters in patients with chronic lymphocytic leukemia and normal controls. *Blut (Berlin)* 50 (1985): 243-248.
- Sipe CR, Chanana AD, Cronkite EP, et al. Studies on lymphocytes XIII. Nuclear volume measurement as a rapid approach to estimate proliferative fraction. *Scand J Haematol* 16 (1976): 196-201.
- National Center for Biotechnology Information (NCBI). *Bos_taurus_UMD_3.1.1* (2016).
- Watanabe S, Kusama-Eguchi K, Kobayashi H, et al. Estimation of polyamine binding to macromolecules and ATP in bovine lymphocytes and rat liver. *J Biol Chem* 266 (1991): 20803-20809.
- National Center for Biotechnology Information (NCBI). *GRCh38.p14* (2022).
- Adey A, Burton JN, Kitzman JO, et al. The haplotype-resolved genome and epigenome of the aneuploid HeLa cancer cell line. *Nature* 500 (2013): 207-211.
- Macville M, Schröck E, Padilla-Nash H, et al. Comprehensive and definitive molecular cytogenetic characterization of HeLa cells by spectral karyotyping. *Cancer Res* 59 (1999): 141-150.
- Goyns MH. Polyamine content of a non-aqueously isolated chromosome preparation. *Exp Cell Res* 122 (1979): 377-380.
- Jain S, Zon G, Sundaralingam M. Base only binding of spermine in the deep groove of the A-DNA octamer d(GTGTACAC). *Biochemistry* 28 (1989): 2360-2364.
- Egli M, Williams LD, Gao Q, et al. Structure of the pure-spermine form of Z-DNA (magnesium free) at 1-Å resolution. *Biochemistry* 30 (1991): 11388-11402.
- Ohishi H, Tozuka Y, Da-Yang Z, et al. The rare crystallographic structure of d(CGCGCG)₂: the natural spermidine molecule bound to the minor groove of left-handed Z-DNA d(CGCGCG)₂ at 10 °C. *Biochem Biophys Res Commun* 358 (2007): 24-28.
- Cameron IL, Smith NKR, Pool TB. Element concentration changes in mitotically active and postmitotic enterocytes. An X-ray microanalysis study. *J Cell Biol* 80 (1979): 444-450.
- National Center for Biotechnology Information (NCBI). *C3H_HeJ_v1* (2016).
- Sarhan S, Seiler N. On the subcellular localization of the polyamines. *Biol Chem Hoppe-Seyler* 370 (1989): 1279-1284.
- Viola-Magni MP, Gahan PB, Pacy J. Phospholipids in plant and animal chromatin. *Cell Biochem Funct* 3 (1985): 71-78.
- Hurst R, Armah CN, Dainty JR, et al. Establishing optimal selenium status: results of a randomized, double-blind, placebo-controlled trial. *Am J Clin Nutr* 91 (2010): 923-931.
- Muecke R, Waldschock K, Schomburg L, et al. Whole blood selenium levels and selenium supplementation in patients treated in a family doctor practice in Golßen (state of Brandenburg, Germany): a laboratory study. *Integr Cancer Ther* 17 (2018): 1132-1136.
- Park YC, Whanger PD. Toxicity, metabolism and absorption of selenite by isolated rat hepatocytes. *Toxicology* 100 (1995): 151-162.
- Calvo SE, Pagliarini DJ, Mootha VK. Upstream open reading frames cause widespread reduction of protein expression and are polymorphic among humans. *Proc Natl Acad Sci USA* 106 (2009): 7507-7512.
- Crowe ML, Wang XQ, Rothnagel JA. Evidence for conservation and selection of upstream open reading frames suggests probable encoding of bioactive peptides. *BMC Genomics* 7 (2006): 16.

28. Kozak M. Compilation and analysis of sequences upstream from the translational start site in eukaryotic mRNAs. *Nucleic Acids Res* 12 (1984): 857-872.
29. Kumler WD, Eiler JJ. The acid strength of mono and diesters of phosphoric acid. The *n*-alkyl esters from methyl to butyl, the esters of biological importance, and the natural guanidine phosphoric acids. *J Am Chem Soc* 65 (1943): 2355-2361.
30. Katchalsky A, Gillis J. Theory of the potentiometric titration of polymeric acids. *Recl Trav Chim Pays-Bas* 68 (1949): 879-897.
31. Kuhn W, Kuhn H. Die Frage nach der Aufrollung von Fadenmolekülen in strömenden Lösungen. *Helv Chim Acta* 26 (1943): 1394-1465.
32. Mandelkern M, Elias JG, Eden D, et al. The dimensions of DNA in solution. *J Mol Biol* 152 (1981): 153-161.
33. Mermer A, Starynowicz P. Charge-density distribution in potassium dihydrogen phosphoglycolate – a comparison of phosphate and phosphonate groups. *Acta Crystallogr B* 68 (2012): 625-635.
34. Aikens D, Bunce S, Onasch F, et al. The interactions between nucleic acids and polyamines. II. Protonation constants and ¹³C-NMR chemical shift assignments of spermidine, spermine, and homologs. *Biophys Chem* 17 (1983): 67-74.
35. Tari LW, Secco AS. Base-pair opening and spermine binding—B-DNA features displayed in the crystal structure of a *gal* operon fragment: implications for protein–DNA recognition. *Nucleic Acids Res* 23 (1995): 2065-2073.
36. Wiberg N (Ed). *Holleman–Wiberg, Lehrbuch der Anorganischen Chemie*, 101. ed, Walter de Gruyter, Berlin, New York (1995): 771.
37. Shor DSM, Struys EA, Hogema BM, et al. Development of a stable-isotope dilution assay for γ -aminobutyric acid (GABA) transaminase in isolated leukocytes and evidence that GABA and β -alanine transaminases are identical. *Clin Chem* 47 (2001): 525–531.
38. Cellini B, Montioli R, Bianconi S, et al. Construction, purification and characterization of untagged human liver alanine-glyoxylate aminotransferase expressed in *Escherichia coli*. *Protein Pept Lett* 15 (2008): 153-159.
39. Oppici E, Fodor K, Paiardini A, et al. Crystal structure of the S187F variant of human liver alanine: glyoxylate aminotransferase associated with primary hyperoxaluria type I and its functional implications. *Proteins* 81 (2013): 1457-1465.
40. Llères D, James J, Swift S, et al. Quantitative analysis of chromatin compaction in living cells using FLIM–FRET. *J Cell Biol* 187 (2009): 481-496.
41. Rhodes CT. Determination of micro-pH in solid drug delivery systems. *Drug Dev Ind Pharm* 25 (1999): 1221-1222.
42. Herbstein FH, Kapon M, Schwotzer W. Crystal structure of tetrakis(phenacetin) dihydrogentetraiodide dihydrate $\{[H_5C_2OC_6H_4N(H)C(CH_3)=O]_4 \cdot H_2I_4 \cdot 2 H_2O\}$. *Helv Chim Acta* 66 (1983): 35-43.
43. Hitchcock PB, Hughes DL, Leigh GJ, et al. Preparation of new vanadium(II) iodides and crystal structure of hexakis(acetonitrile)vanadium(II) (tetraiodide). *J Chem Soc, Dalton Trans* (1994): 3683-3687.
44. Genser EE, Connick RE. Exchange of iodide ion with triiodide ion studied by nuclear magnetic resonance. *J Chem Phys* 58 (1973): 990-996.
45. Schenck H-U, Simak P, Haedicke E. Structure of polyvinylpyrrolidone-iodine (povidone-iodine). *J Pharm Sci* 68 (1979): 1505-1509.
46. Herbstein FH, Kapon M. Zigzag chains of alternating iodine molecules and triiodide ions in crystalline (phenacetin)₂ • HI₅. *Nature Phys Sci* 239 (1972): 153-154.
47. Saenger W. The structure of the blue starch–iodine complex. *Naturwissenschaften* 71 (1984): 31-36.
48. Svensson PH, Kloo L. Synthesis, structure, and bonding in polyiodide and metal iodide–iodine systems. *Chem Rev* 103 (2003): 1649-1684.
49. Tomlin CDS (Ed). *The Pesticide Manual: A World Compendium, Incorporating the Agrochemicals Handbook*, eleventh ed, British Crop Protection Council, Farnham, Surrey, UK (1997).
50. Terpstra KR, Woortman AJJ, Hopman JCP. Yellow dextrans: evaluating changes in structure and colour during processing. *Starch/Stärke* 62 (2010): 449-457.
51. Cellini B, Bertoldi M, Montioli R, et al. Human wild-type alanine:glyoxylate aminotransferase and its naturally occurring G82E variant: functional properties and physiological implications. *Biochem J* 408 (2007): 39-50.
52. Schuster H, Schramm G, Zillig W. Die Struktur der Ribonucleinsäure aus Tabakmosaikvirus. *Z Naturforsch B* 11 (1956): 339-345.
53. Takeda Y, Samejima K, Nagano K, et al. Determination of protonation sites in thermospermine and in some other polyamines by ¹⁵N and ¹³C nuclear magnetic resonance spectroscopy. *Eur J Biochem* 130 (1983): 383-389.
54. Waddington CH. *The epigenotype*. Endeavour (Engl Ed London) 1 (1942): 18-20.

55. Crick FHC, Watson JD. The complementary structure of deoxyribonucleic acid. *Proc R Soc Lond A Math Phys Sci* 223 (1954): 80-96.
56. Saenger W, Hunter WN, Kennard O. DNA conformation is determined by economics in the hydration of phosphate groups. *Nature* 324 (1986): 385-388.
57. Sunkara PS, Ramakrishna S, Nishioka K, et al. The relationship between levels and rates of synthesis of polyamines during mammalian cell cycle. *Life Sci* 28 (1981): 1497-1506.
58. Li G, Sudlow G, Belmont AS. Interphase cell cycle dynamics of a late-replicating, heterochromatic homogeneously staining region: precise choreography of condensation/decondensation and nuclear positioning. *J Cell Biol* 140 (1998): 975-989.
59. Flohé L, Andreesen JR, Brigelius-Flohé R, et al. Selenium, the element of the moon, in life on earth. *IUBMB Life* 49 (2000): 411-420.
60. Berzelius JJ. Lettre de M. Berzelius à M. Berthollet sur deux Métaux nouveaux. *Ann Chim Phys (Paris)* 7 (1817): 199-206.
61. Borah S, Kumar PP. *Ab initio* molecular dynamics study of Se (IV) species in aqueous environment. *Phys Chem Chem Phys* 18 (2016): 26755-26763.
62. Nafisi S, Montazeri M, Manouchehri F. The effect of Se salts on DNA structure. *J Photochem Photobiol B* 113 (2012): 36-41.
63. Nafisi S, Manouchehri F, Montazeri M. RNA adducts with Na₂SeO₄ and Na₂SeO₃ – Stability and structural features. *J Mol Struct* 1006 (2011): 547-552.
64. Rueppel ML, Brightwell BB, Schaefer J, et al. Metabolism and degradation of glyphosate in soil and water. *J Agric Food Chem* 25 (1977): 517-528.
65. Benbrook CM. Trends in glyphosate herbicide use in the United States and globally. *Environ Sci Eur* 28 (2016): 3.
66. Holmes RP, Assimos DG. Glyoxylate synthesis, and its modulation and influence on oxalate synthesis. *J Urol* 160 (1998): 1617-1624.
67. Helgstrand E, Eriksson B, Johansson NG, et al. Trisodium phosphonoformate, a new antiviral compound. *Science* 201 (1978): 819-821.
68. Sabourin CLK, Reno JM, Boezi JA. Inhibition of eucaryotic DNA polymerases by phosphonoacetate and phosphonoformate. *Arch Biochem Biophys* 187 (1978): 96-101.
69. Khan S, Ahmed S. Role of *swi7H4* mutant allele of DNA polymerase α in the DNA damage checkpoint response. *PLoS One* 10 (2015): e0124063.
70. Rosa S, Shaw P. Insights into chromatin structure and dynamics in plants. *Biology* 2 (2013): 1378-1410.
71. Zeng H. Selenium as an essential micronutrient: roles in cell cycle and apoptosis. *Molecules* 14 (2009): 1263-1278.
72. Frenkel GD. Effects of sodium selenite and selenate on DNA and RNA synthesis in vitro. *Toxicol Lett* 25 (1985): 219-223.
73. Loeb LA, Preston BD. Mutagenesis by apurinic/aprimidinic sites. *Ann Rev Genet* 20 (1986): 201-230.
74. Miller CO, Skoog F, Okumura FS, et al. Isolation, structure and synthesis of kinetin, a substance promoting cell division. *J Am Chem Soc* 78 (1956): 1375-1380.
75. Barciszewski J, Siboska GE, Pedersen BO, et al. A mechanism for the in vivo formation of N⁶-furfuryladenine, kinetin, as a secondary oxidative damage product of DNA. *FEBS Lett* 414 (1997): 457-460.
76. Barciszewski J, Massino F, Clark BFC. Kinetin—a multiactive molecule. *Int J Biol Macromol* 40 (2007): 182-192.
77. Fuller WD, Sanchez RA, Orgel LE. Studies in prebiotic synthesis. VI. Synthesis of purine nucleosides. *J Mol Biol* 67 (1972): 25-33.
78. Maurel MC, Convert O. Chemical structure of a prebiotic analog of adenosine. *Orig Life Evol Biosph* 20 (1990): 43-48.
79. Mustard TJL, Mack DJ, Njardarson JT, et al. Mechanism and the origins of stereospecificity in copper-catalyzed ring expansion of vinyl oxiranes: a traceless dual transition-metal-mediated process. *J Am Chem Soc* 135 (2013): 1471-1475.
80. Hardi EA, Njardarson JT. Ring expansions of vinyloxiranes, -thiiranes, and -aziridines: synthetic approaches, challenges, and catalytic success stories. *J Org Chem* 78 (2013): 9533-9540.
81. de Alencastro G, McCloskey DE, Kliemann SE, et al. New *SMS* mutation leads to a striking reduction in spermine synthase protein function and a severe form of Snyder–Robinson X-linked recessive mental retardation syndrome. *J Med Genet* 45 (2008): 539-543.
82. Koller VJ, Fürhacker M, Nersesyan A, et al. Cytotoxic and DNA-damaging properties of glyphosate and Roundup in human-derived buccal epithelial cells. *Arch Toxicol* 86 (2012): 805-813.
83. Coupe RH, Kalkhoff SJ, Capel PD, et al. Fate and transport of glyphosate and aminomethylphosphonic acid in surface waters of agricultural basins. *Pest Manag Sci* 68 (2012): 16-30.